

Fisheries Long Term Monitoring Program

Syngnathids and their associated communities

Supplementary report to:
Syngnathids in the East Coast Trawl Fishery: a review and trawl survey

August 2006



Fisheries Long Term Monitoring Program

Syngnathids and their associated communities

Supplementary report to:

Syngnathids in the East Coast Trawl Fishery: a review and trawl survey

August 2006

Natalie Dodt

Department of Primary Industries and Fisheries

Queensland

ISSN 0727-6273

QI06096

This document may be cited as:

N. Dodt (2006). Fisheries Long Term Monitoring Program - Syngnathids and their associated communities - Supplementary report to: Syngnathids in the East Coast Trawl Fishery: a review and trawl survey. Department of Primary Industries and Fisheries, Brisbane, Australia.

Acknowledgements:

Thanks are due to the commercial fishermen John and Gavin McIlwain for their willingness to undertake the survey work. I would like to acknowledge the Long Term Monitoring Program team members and fisheries observers for collecting the samples at sea and processing them in the laboratory. I am grateful to Malcolm Dunning, Eddie Jebreen and Olivia Whybird, all of whom reviewed previous versions of this report.

Particular thanks must go to Jonathan Staunton-Smith and Darren Roy for their support and assistance in this project.

The Department of Primary Industries and Fisheries (DPI&F) seeks to maximise the economic potential of Queensland's primary industries on a sustainable basis.

This publication has been compiled by Fisheries, Assessment and Monitoring.

While every care has been taken in preparing this publication, the State of Queensland accepts no responsibility for decisions or actions taken as a result of any data, information, statement or advice, expressed or implied, contained in this report.

© The State of Queensland, Department of Primary Industries and Fisheries 2006

Copyright protects this material. Except as permitted by the *Copyright Act 1968* (Cth), reproduction by any means (photocopying, electronic, mechanical, recording or otherwise), making available online, electronic transmission or other publication of this material is prohibited without the prior written permission of The Department of Primary Industries and Fisheries, Queensland. Inquiries should be addressed to copyright@dpi.qld.gov.au (telephone +61 7 3404 6999).

Inquiries should be addressed to:

Intellectual Property and Commercialisation Unit

Department of Primary Industries and Fisheries

GPO Box 46

Brisbane Qld 4001

Contents

Contents	iii
Acronyms	iii
Introduction	1
Methods	1
Data Analysis	3
Results	5
Discussion	10
References	11
Appendix 1	12

Acronyms

BRD	Bycatch Reduction Device
CRW	Catch Rate - Weight
CRN	Catch Rate - Numbers
MDS	Multidimensional Scaling
PRIMER	Plymouth Routines in Multivariate Ecological Research
SIMPER	Similarity of Percentages
TED	Turtle Exclusion Device

Introduction

This document presents the data collected to address the fourth objective of the initial report (Dodt 2005), to “investigate the relationship between syngnathid distribution and abundance, and assemblages and habitat characteristics”.

Methods

Dodt (2005) details the survey design, field sampling protocols and laboratory procedures for collection and processing of the syngnathid samples from the trawl survey. The location of the survey shots and the stratified grids sampled during the survey are shown in Figure 1. From each trawl shot a 10 kg sub-sample of the catch was collected from the middle net to evaluate community composition¹. The latitude and longitude, depth and strata information for each shot was recorded. Large individuals (including sharks, rays, sponges and habitat characters (i.e. rubble)) caught in the middle net that could not be accurately sub-sampled, due to their size, were identified to the lowest taxonomic level achievable, number of individuals and their weight were recorded on board the vessel and released. These individuals are referred to as “monsters” in this report. Syngnathids were removed from the catch prior to collecting the sub-sample. The details of syngnathid processing are in the initial report (Dodt 2005).

In order to calculate catch rates of the species and habitat characters in each sub-sample the following information was collected from each trawl:

- total weight of catch in the middle net (full net weight - empty net weight)
- wet weight of the sub-sample (sum of weights of all the contents as processed in the laboratory)
- total weight of each species (including habitat characters) in the sub-sample
- number of individuals of each species in the sub-sample.

The sub-sample from each trawl shot was sorted to lowest taxonomic level. Species² were identified according to protocols outlined in the Department of Primary Industries and Fisheries species register (Roy *et al.* 2005).

The lengths of up to 20 randomly selected individuals were recorded for each species in each sub-sample. Only maximum and minimum sizes are presented in this report.

¹ *Community Composition* refers to all catch (including rubble or sediment, yet excluding syngnathids for this survey) trawled by nets without excluder devices such as TEDs and BRDs.
² In order to be consistent with terminology throughout this section, we use the term species (plural form) even when referring to groupings at the higher taxonomic level

