# QUEENSLAND DEPARTMENT OF PRIMARY INDUSTRIES

ì

DIVISION OF PLANT INDUSTRY BULLETIN No. 366

# THE KANGAROO INDUSTRY IN QUEENSLAND. FIRST SUPPLEMENT, 1961–1965

By C. ROFF AND T. KIRKPATRICK, M.Sc.

#### SUMMARY

The total harvest for the period 1961-1965 was nearly four million marsupials, an increase of some 25% on that of the previous 5 years. Practically all this increase involved the grey kangaroo: from 1,450,000 to 2,490,000.

Estimated annual income of the industry at the level of prices to shooters increased from £150,000 (1950-1960) to £450,000 (skins £390,000, carcasses £60,000) (1961-65).

Information on the kangaroo industry in Queensland before and including 1960 was presented by Roff and Kirkpatrick (1962). During the past 5 years there has been a distinct improvement in field recording at the species level, and the use of the category unspecified was not required. For continuity and convenience some of the earlier information (Roff and Kirkpatrick 1962) has been brought forward either as actual figures or yearly averages, and included in the tables for 1961-1965.

"Queensland Journal of Agricultural and Animal Sciences", Vol. 23, 1966

## SHORTER COMMUNICATIONS

## TABLE 1

1

ð

Ł

# NUMBERS OF MARSUPIAL SKINS HARVESTED, 1954–1960 and 1961–1965

	Total	Number and Percentage by Species									
Year	Number Harvested	Grey Kangaroo	Red Kangaroo	Wallaroo	Whiptail Wallaby	Brush Wallaby	Swamp Wallaby	Sandy Wallaby			
1954	218,459	127,753 (58·47)	68,683 (31·43)	3,295 (1·50)	6,429 (3·61)	7,907 (2·94)	1,678 (0·76)	2,714 (1·24)			
1955	305,616	155,956 (51·03)	93,616 (30·63)	11,155 (3·65)	11,580 (3·80)	15,561 (5·09)	12,848 (4·20)	4,900 (1·60)			
1956	361,462	183,663 (50·81)	111,034 (30·71)	21,642 (5•98)	13,800 (3·81)	13,923 (3·85)	12,167 (3·36)	5,233 (1·44)			
1957	631,034	283,137 (44·86)	276,125 (43·75)	29,897 (4·73)	13,781 (2·18)	14,756 (2·33)	6,959 (1·10)	6,379 (1·01)			
1958	295,820	172,437 (58·29)	89,248 (30·16)	12,212 (4·12)	11,164 (3·77)	7,063 (2·38)	3,168 (1·07)	528 (0·17)			
1959	1,006,919	319,829 (31·76)	588,807 (58·47)	55,031 (5·46)	9,173 (0·91)	25,141 (2·49)	6,784 (0·67)	2,154 (0·21)			
1960	769,948	488,118 (63·39)	250,059 (32·47)	10,319 (1·34)	5,478 (0·71)	8,059 (1·04)	6,642 (0·86)	1,273 (0·16)			
1961	471,640	293,365 (62·20)	161,063 (34·14)	3,374 (0·71)	11,038 (2·34)	2,208 (0·46)	592 (0·12)	Nil			
1962	548,771	369,631 (67·35)	163,712 (29·83)	8,377 (1·52)	4,945 (0·90)	1,227 (0·22)	879 (0·16)	Nil			
1963	642,063	444,607 (69·24)	175,284 (27·30)	6,727 (1·04)	11,026 (1·71)	3,462 (0·53)	846 (0·13)	111 (0·01)			
1964	1,118,813	596,775 (53·34)	496,439 (44·37)	13,328 (1·19)	7,800 (0·69)	3,299 (0·29)	1,172 (0·10)	Nil			
1965	1,168,887	787,955 (67·41)	353,223 (30·21)	13,365 (1·14)	8,662 (0·74)	4,139 (0·35)	1,455 (0·12)	88 (0·007			

468

Species	Year	Peninsula	Cairns	Towns- ville	Mackay	Rock- hampton	Mary- borough	Moreton	Downs	Roma	South Western	Far Western	Central Western	North Western
Grey kangaroo	1961 1965	···	214 2	270 2,104	200 46	62,996 23,636	10,707 14,938	10,390 10,171	39,262 165,940	135,747 439,009	22,112 36,622	 900	11,453 84,073	14 10,514
Red kangaroo	1961 1965		· · ·	 162	· · ·	3,995	2,137 1,630	3,422 391	5,202 27,989	23,595 34,815	89,483 146,888	1,504 27,241	31,256 73,202	4,464 36,910
Wallaroo	1961 1965			34 157		5 1,131	196 4	227 94	1,432 1,860	97 432	503 1,611	250 395	506 5,827	124 1,826
Whiptail wallaby	1961 1965		259 1	41 25	304 17	1,542 3,798	5,519 2,029	2,007 802	1,263 1,351	46 615			4 24	53
Brush wallaby	1961 1965		8	5	30 14	219 676	141 405	227 438	1,448 1,474	78 762	 107	 19	4 24	53 215
Swamp wallaby	1961 1965		•••		133 20	125 111	10 235	22 12	215 634	44 158	43 11		 274	
Sandy wallaby	1961 1965	· · ·				· · · · · ·	· · · · · ·	  						· · · · · · · · · · · · · · · · · · ·
All species	1954 and 1959	15*	61*	625*	227*	15,367*	7,229*	3,499*	61,212*	86,162*	285,701*	2,527*	69,168*	7,487*
All species	1961 and 1965		255*	1,443*	383*	49,117*	18,976*	14,102*	124,035*	317,699*	148,690*	15,155*	103,324*	27,087*

## TABLE 2

Śm

GEOGRAPHIC ORIGIN OF SKINS: HARVESTS IN STATISTICAL DIVISIONS DURING 1954 AND 1959, 1961 AND 1965

\* Yearly average.

SHORTER COMMUNICATIONS

~ \_\_\_\_\_

s ~

469

## TABLE 3

## NUMBERS OF SANDY AND WHIPTAIL WALLABIES TAKEN IN QUEENSLAND, 1956–1960 AND 1961–1965

15

4

ſ.

Period/Year	Number Taken as Pests in Sugar-cane Districts*	Number Harvested Elsewhere and Skins Marketed
1956–1960	61,287	68,737
1961 1962 1963	18,826 14,172 14,143	11,038 4,945 11,137
1964 1965	10,418 11,468	7,800 8,750
1961–1965	69,027	43,670

\* Numbers on which a bounty was paid.

### TABLE 4

#### ESTIMATED WEIGHT OF KANGAROO MEAT PRODUCED IN QUEENSLAND AND NUMBERS OF CARCASSES, 1959–1960 and 1961–1965

Year	Weight (long tons)	Number of Carcasses	Percentage of Total Kangaroo Harvest		
1959	39.00	2,500	0.27		
1960	824.96	67,100	9.08		
1961*	850.00	85,000	18.70		
1962	971.30	97,130	18.21		
1963	827.20	82,720	13.34		
1964	2,304.13	230,413	21.07		
1965	4,068.71	406,871	35.65		

\* Information given for 1961 in Table 4 of Roff and Kirkpatrick (1962) was for part year only.

Period/Year	Peninsula	Cairns	Towns- ville	Mackay	Rock- hampton	Mary- borough	Moreton	Downs	Roma	South Western	Far Western	Central Western	North Western
1956–1960		7*	17*	4*	204*	64*	90*	472*	359*	273*	36*	221*	57*
1961		3	3	2	140	85	22	444	279	284	13	175	20
1962		1	10		204	86	40	248	275	307	29	298	16
1963		1	14	4	227	99	143	398	289	218	14	264	40
1964			7	6	242	80	31	513	396	300	49	496	78
1965		• •	6	1	168	59	26	525	500	407	38	350	73
1961–1965		1*	8*	3*	196*	82*	52*	426*	348*	303*	29*	317*	45*

TABLE 5

DISTRIBUTION OF REGISTERED SHOOTERS IN STATISTICAL DIVISIONS, 1956–1960 and 1961–1965

Ę.,

)5

-

\* Yearly average.

471

~ »-

-là

4

TABLE	6
-------	---

BRISBANE PRICES FOR KANGAROO SKINS (PER LB), 1961–1965\*

1

A.

e li

						,,	
Month		Grade	1961	1962	1963	1964	1965
May	• •	1 2 3	s. d. 4 6 3 4 1 9	s. d. 5 6 4 0 3 0	s. d. 4 6 3 6 3 0	s. d. 6 6 5 9 4 6	s. d. 5 9 4 3 3 0
June		1 2 3	6 3 4 7 2 9	6 8 5 6 3 4	7 6 6 6 6 0	7 0 6 0 4 6	6 9 5 0 3 0
July	•••	1 2 3	6 3 4 7 2 9	6 11 5 9 3 6	8 6 7 6 7 0	7 0 6 0 4 6	6 9 5 0 3 0
August		1 2 3	6 4 4 9 2 9	6 9 5 8 3 6	9 0 8 0 7 0	†6 10 †6 0 †4 0	6 9 5 0 3 0
September		1 2 3	6 0 4 11 3 0	6 9 5 8 3 6	8 10 7 10 6 11	6 6 5 0 3 6	6 9 5 0 3 0
October		1 2 3	4 9 3 9 3 0	69 56 40	.8 4 7 4 6 6	6 9 5 0 3 6	6 9 5 0 3 0

\* An average skin weighs  $1-1\frac{3}{4}$  lb.

† Some skins unsaleable late August.

TABLE 7

DATA CONCERNED WITH FLUCTUATIONS OF ANNUAL HARVESTS, 1956–1960 AND 1961–1965

Period/Year	Number of Registered Shooters	Number of Kangaroos Marketed for Skins and Meat	Seasonal Conditions	for 1st G	1 Average rade Skins r lb)	Estimated Average Price for Meat (per lb)	
1956–1960	1,805*	552,405*†		<i>s</i> . 6	<i>d.</i> 8	<i>d</i> .	
1961	1,470	454,428	Fair	5	8	3.5	
1962	1,514	533,343	Fair	6	6	3.5	
1963	1,711	619,891	Fair	7	9	3.5	
1964	2,198	1,093,214	Dry	6	8	3.5	
1965	2,155	1,141,178	Drought	6	7	3.5	
1961–1965	1,809*	768,410*		6	8*	3.5*	

\* Yearly average.

<sup>†</sup> Number of kangaroos marketed in 1956 is given as 194,663 (Roff and Kirkpatrick 1962). Corrected figure of 294,663 used.

#### SHORTER COMMUNICATIONS

#### Comment

"During dry seasons kangaroos are more readily and economically available, as the mobs congregate near the limited water and food; furthermore, more rural workers turn to skin shooting to supplement incomes" (Roff and Kirkpatrick 1962). This seasonal influence on harvest was emphasized again in the data for 1961-1965. During the fair-season years of 1961-1963 the annual take of kangaroos was of the order of half a million, while in the dry and drought years of 1964-1965 this was doubled: in parallel, the number of registered shooters increased by some 39% (Tables 1 and 7).

The total harvest for the period 1961-1965 was approximately four million marsupials, an increase of about 25% on that of the previous 5 years. Nearly all this increase involved the grey kangaroo: from 1,450,000 to 2,490,000. The 5-year kills of red kangaroos were respectively 1,315,000 and 1,350,000 (Table 1).

There has been an increase in shooting in the middle districts of the State i.e. west of the Great Dividing Range but excluding the far western areas. This increase is most obvious in the more southern of these middle districts (Table 2).

The marketing of carcasses (meat industry) has expanded, with a record sale of some 4,000 long tons in the drought year of 1965 (Table 4).

Estimated annual income of the kangaroo industry at the level of prices to shooters has increased from £150,000 during 1950-1960 (Roff and Kirkpatrick 1962) to £450,000 (skins £390,000, carcasses £60,000) for the period 1961-1965.

The effects of harvesting on kangaroo populations and conservation, and the meat industry, will be discussed in detail elsewhere.

### REFERENCE

ROFF, C., and KIRKPATRICK, T. (1962).—The kangaroo industry in Queensland. Qd J. Agric. Sci. 19:385-401.

#### (Received for publication August 31, 1966)

The authors are officers of the Entomology Section, Division of Plant Industry, Department of Primary Industries.

S. G. REID, Government Printer, Brisbane