

A systematic review of ground-based shooting to control overabundant mammal populations

Andrew J. Bengsen^{A,E}, David M. Forsyth^A, Stephen Harris^B, A. David M. Latham^C, Steven R. McLeod^A and Anthony Pople^D

^AVertebrate Pest Research Unit, NSW Department of Primary Industries, 1447 Forest Road, Orange, NSW 2800, Australia.

^BSchool of Biological Sciences, Life Sciences Building, 24 Tyndall Avenue, Bristol BS8 1TQ, United Kingdom.

^CWildlife Ecology and Management, Manaaki Whenua – Landcare Research, PO Box 69040, Lincoln 7640, New Zealand.

^DInvasive Plants and Animals, Biosecurity Queensland, Department of Agriculture and Fisheries, 41 Boggo Road, Dutton Park, Qld 4102, Australia.

^ECorresponding author. Email: andrew.bengsen@dpi.nsw.gov.au

Table S1. Summary of studies used in the systematic review of ground-based shooting to control overabundant mammal populations, arranged by taxonomic Order, Family and Genus

| Ref | Species | Main shooters | Objective | Target metric | Region | Landscape | Status | Population change indicator | Damage indicator | Main conclusions relevant to this review |
|-----|-------------------------------|---------------------|-------------|---------------|---------------|-----------|--------|-----------------------------|-------------------|---|
| 1 | <i>Capra hircus</i> | Government | None | Density | Australasia | Mixed | Exotic | Relative abundance | na | Goats were nearly eradicated from a small area, but the population recovered substantially within a year due to immigration and survivors' reproduction. |
| 2 | <i>Capra hircus</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Abundance | na | Hunting with dogs reduced and suppressed the goat population over 38 years. This resulted in a visible, but not measured, recovery of susceptible plant spp. Eradication would require annual removal of > 90% of the population. |
| 3 | <i>Capra hircus</i> | Government | Eradication | Eradication | Australasia | Mixed | Exotic | Presence | Native vegetation | A prolonged shooting program using a small number of staff successfully eradicated feral goats. |
| 4 | <i>Capra hircus</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Presence | na | Eradication was unsuccessful, despite high efficiency, because of reinvasion from adjacent areas that were not targeted. |
| 4 | <i>Capra hircus</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Presence | na | Eradication, mainly by ground hunting with dogs, was successful but inefficient. |
| 5 | <i>Hemitragus jemlahicus</i> | Recreational hunter | Density | Density | Australasia | Native | Exotic | Abundance; Class Structure | na | Recreational hunting did not reduce tahr densities below the 'intervention threshold' of 2.5 tahr km ⁻² . Tahr increased over 12 years of study at two sites, despite hunting. |
| 6 | <i>Ovis aries, O. gmelini</i> | Recreational hunter | Eradication | Eradication | North America | Native | Exotic | Presence | Native vegetation | 21 years of public hunting and supplementary aerial shooting failed to eradicate or reduce ungulate (mainly sheep) numbers and impacts. Threatened bird (palila) is declining, but also due to factors other than introduced ungulates. |
| 7 | <i>Capreolus capreolus</i> | Recreational hunter | None | Damage | Europe | Native | Native | na | Native vegetation | Browse damage decreased, on average, across most game management districts where management recommendations were made to increase the harvest. Damage decrease was greater than expected from reduced deer density |

| | | | | | | | | | | |
|----|----------------------------|---|-------------|-------------|-------------|--------|--------|--------------------|-------------------|--|
| | | | | | | | | | | alone. |
| 8 | <i>Capreolus capreolus</i> | Recreational hunter | None | Density | Europe | Mixed | Native | Density | na | Forest and grassland areas were sources of emigrants, despite active management from three types of shooters. This was attributed to the absence of informed management objectives, rather than a failure of shooting operations to meet objectives. |
| 9 | <i>Cervus elaphus</i> | Commercial wildlife management contractor | Eradication | Eradication | Australasia | Native | Exotic | Presence | na | Secretary Island: five year eradication program removed 633 deer: 542 during initial knockdown phase (aiming for 80%) and 91 during two year mop up. Mop up was ongoing when the paper was published. |
| 10 | <i>Cervus elaphus</i> | Government | None | Density | Australasia | Native | Exotic | Abundance | na | Waihaha site only (only site with significant ground hunting): index declined in nil-treatment site, but not at ground hunting site. Helicopter hunting was more effective at sites where it could be used. |
| 11 | <i>Cervus elaphus</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Relative abundance | na | Deer populations were stable over the four years of the study. Hunters probably contributed substantially to inhibiting population growth. |
| 12 | <i>Cervus elaphus</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Relative abundance | Native vegetation | Recreational hunting had some benefit in maintaining an intact forest canopy, but much lower deer densities would be required to reverse the damage that has been caused by deer. |
| 13 | <i>Cervus elaphus</i> | Recreational hunter | Density | Density | Europe | Native | Native | Density | na | Structured population model from bag records and genetic capture-mark-recapture both indicated a decrease at the site over 10 years, but not yet to desired levels, while bag records increased at other sites. |
| 14 | <i>Cervus elaphus</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Relative abundance | Native vegetation | Aerial shooting was effective at reducing deer and increasing seedling growth in the high intensity site, but recreational and commercial shooting had no impact on deer density or seedling growth in the other 3 sites. |

| | | | | | | | | | | |
|----|------------------------|---|-------------|-------------|-------------|--------|--------|--------------------|-------------------|--|
| 9 | <i>Cervus elaphus</i> | Commercial wildlife management contractor | Eradication | Eradication | Australasia | Native | Exotic | Presence | na | Anchor Island: five year eradication program removed 29 deer and no deer were found in the following two years. Almost all deer were killed by ground hunters because conditions were unsuitable for baiting or aerial shooting. |
| 10 | <i>Cervus nippon</i> | Government | None | Density | Australasia | Native | Exotic | Abundance | na | Waihaha site only (only site with significant ground hunting): Index declined in nil-treatment site, but not at ground hunting site. Helicopter hunting was more effective at sites where it could be used. |
| 15 | <i>Cervus nippon</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Abundance | na | Recreational hunting slowed the rate of population growth, but the population only began to decline after very intensive government control started. |
| 12 | <i>Cervus nippon</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Relative abundance | Native vegetation | Recreational hunting had some benefit in maintaining an intact forest canopy, but much lower deer densities would be required to reverse the damage that has been caused by deer. |
| 14 | <i>Cervus nippon</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Relative abundance | Native vegetation | Aerial shooting was effective at reducing deer and increasing seedling growth in the high intensity site, but recreational and commercial shooting had no impact on deer density or seedling growth in the other three sites. |
| 16 | <i>Cervus nippon</i> | Recreational hunter | Growth | Growth | Asia | Native | Native | Modelled Abundance | na | Emergency population control (maximise female harvest) measures initiated after high estimated densities failed to reduce the population to desired levels. |
| 17 | <i>Cervus nippon</i> | Recreational hunter | None | Density | Asia | Native | Native | Relative abundance | na | Effects of localised hunting and culling were outweighed by movement between sites. |
| 18 | <i>Cervus unicolor</i> | Commercial wildlife management contractor | None | Damage | Australasia | Native | Exotic | Relative abundance | Faecal density | Intensive bouts of ground shooting reduced faecal deposition and pollution risk near the water, but the effects only persisted while shooting was occurring. |

| | | | | | | | | | | |
|----|-------------------------------|---|-------------|-------------|---------------|------------|--------|--------------------|--------------------|--|
| 19 | <i>Dama dama</i> | Government | Eradication | Eradication | Australasia | Mixed | Exotic | Presence | na | Evaluation of 54 programs aiming to eradicate new deer populations, mainly using ground based hunting. 35/54 programs were effective, and the overall program was thought to be effective. |
| 3 | <i>Dama dama</i> | Government | Eradication | Eradication | Australasia | Mixed | Exotic | Presence | na | A prolonged shooting program using a small number of staff successfully eradicated deer. |
| 20 | <i>Dama dama</i> | Recreational hunter | None | Density | Australasia | Native | Exotic | Density | na | The reliability of recreational hunters to control deer populations depended on the size of the hunter population within 100 km and on the extent of road access. |
| 8 | <i>Muntiacus reevesi</i> | Recreational hunter | None | Density | Europe | Mixed | Exotic | Density | na | Forest and grassland areas were sources of emigrants, despite active management from three types of shooters. This was attributed to the absence of informed management objectives, rather than a failure of shooting operations to meet objectives. |
| 21 | <i>Odocoileus hemionus</i> | Recreational hunter | None | Damage | North America | Native | Exotic | na | Native vegetation | Deer impacts on tree regeneration were lower in sites with good hunter access, although hunting pressure was probably only about 10% of mortality. |
| 22 | <i>Odocoileus virginianus</i> | Commercial wildlife management contractor | None | Density | North America | Peri urban | Native | Density | Vehicle collisions | Targeted shooting reduced deer populations and deer:vehicle collisions by an average of 67% at three suburban sites. |
| 23 | <i>Odocoileus virginianus</i> | Government | None | Density | North America | Peri urban | Native | Relative abundance | Vehicle collisions | Public hunting combined with three different types of government shooting reduced deer density index by 46% and reported vehicle collisions by 30%. |
| 24 | <i>Odocoileus virginianus</i> | Government | Growth | Growth | North America | Mixed | Native | Abundance | na | Shooting effectively reduced deer densities from very high (136 km ²) to acceptable (<10 km ²). Maintenance at that level was thought to be achievable and affordable (however, a later study showed higher densities and concluded it was impossible to reach 10 km ²). |
| 25 | <i>Odocoileus virginianus</i> | Recreational hunter | None | Density | North America | Peri urban | Native | Relative abundance | Native vegetation | A very intensively managed 15 year hunting program reduced deer density to acceptable levels within a small peri-urban reserve and adjacent agricultural area, while deer density and damage |

| | | | | | | | | | | |
|----|-------------------------------|---|-------------|-------------|---------------|------------|--------|--------------------|--------------------|---|
| | | | | | | | | | | continued to increase in adjacent suburbs. |
| 26 | <i>Odocoileus virginianus</i> | Recreational hunter | Density | Density | North America | Native | Native | Density | na | Recreational hunting reduced deer densities to desired levels within a few years at all blocks within the broader area. |
| 27 | <i>Odocoileus virginianus</i> | Recreational hunter | Growth | Growth | North America | Native | Native | Relative abundance | Native vegetation | Antlerless deer harvest objectives were achieved, but the program wasn't considered effective because it didn't translate into reduced deer density or reduced damage to vegetation. |
| 28 | <i>Odocoileus virginianus</i> | Volunteer | Density | Density | North America | Peri urban | Native | Density | Vehicle collisions | Public and private land volunteer shooting greatly reduced deer densities at four sites, but did not reach management objective of 10 deer km ⁻² . |
| 29 | <i>Sus scrofa</i> | Commercial wildlife management contractor | Eradication | Eradication | North America | Native | Exotic | Abundance | na | Professional hunting was effective at reducing abundance to near zero at most sites, but its effects were compromised by immigration. |
| 30 | <i>Sus scrofa</i> | Volunteer | None | Growth | Australasia | Native | Exotic | Mortality | na | Hunting removed 22/79 pigs. Authors concluded it was useful after other control methods had been exhausted. |
| 31 | <i>Sus scrofa</i> | Recreational hunter | None | Density | North America | Native | Exotic | Relative abundance | na | Hunting, combined with trapping, reduced pig activity within a couple of years before stabilising at a lower level. Hunting needed to be more spatially and temporally widespread to have a greater impact. |
| 32 | <i>Sus scrofa</i> | Recreational hunter | None | Damage | Europe | Mixed | Native | na | Crop damage | Crop damage was negatively, but weakly, related to increasing hunting activity. Fencing and supplementary feeding had negligible benefit. |
| 33 | <i>Sus scrofa</i> | Commercial harvest | Growth | Growth | Australasia | Mixed | Exotic | Abundance | na | Commercial harvesting was incapable of suppressing pig populations. Harvest rates were generally <30% of the estimated population size; much less than the minimum consistent harvest rate that would be expected to inhibit population growth. |
| 34 | <i>Sus scrofa</i> | Recreational hunter | Damage | Damage | South America | Native | Exotic | Relative abundance | Rooting | Sustained hunting effort reduced relative abundance of pigs and maintained at low levels for 10 years. There was a rapid effect on pig rooting and damage to |

| | | | | | | | | | | |
|----|----------------------|---|-------------|-------------|---------------|------------|--------|--------------------|---------------------|---|
| | | | | | | | | | | palms, but it took four years to achieve the objective level. |
| 35 | <i>Sus scrofa</i> | Recreational hunter | None | Growth | North America | Native | Exotic | Growth, Survival | na | Hunting reduced survival but not population growth. |
| 36 | <i>Sus scrofa</i> | Recreational hunter | None | Growth | Europe | Mixed | Native | Survival | na | Harvest rate of recreational hunters was < total net reproduction. |
| 37 | <i>Sus scrofa</i> | Government | None | Density | Australasia | Native | Exotic | Density | Diggings | Intensive control reduced estimated pig density by over one third and this resulted in a decrease in rooting damage >50%. Modelling of different culling intervals indicated that only the most intense (3 monthly intervals) would maintain low damage levels. |
| 38 | <i>Sus scrofa</i> | Government | None | Growth | Australasia | Native | Exotic | Survival | na | Hunting with dogs could be useful for controlling pigs in forested areas, but was likely to be more costly and less effective than poison baiting. |
| 39 | <i>Sus scrofa</i> | Commercial wildlife management contractor | Eradication | Eradication | North America | Native | Exotic | Presence | na | Ground hunting was critical for killing the last pigs. Fixed price funding model for commercial contractor contributed to efficient eradication. |
| 40 | <i>Sus scrofa</i> | Recreational hunter | None | Growth | Europe | Native | Native | Growth | na | Population growth was greatest after six short term (1-3 year) hunting bans were relaxed, suggesting that hunting helped to slow population growth. However, negative population growth was uncommon and inconsistent across sites and years. |
| 41 | <i>Sus scrofa</i> | Government | None | Growth | Europe | Native | Native | Growth | na | Positive population growth in two lightly and heavily hunted populations, but there were differences in many demographic parameters. Harvest rates need to increase substantially to suppress growth. |
| 42 | <i>Sus scrofa</i> | Recreational hunter | None | Growth | Europe | Native | Native | Survival | na | Annual, seasonal recreational drive hunting did not prevent 5-fold population growth over 20 years despite high hunting and total mortality (>50%). |
| 43 | <i>Vulpes vulpes</i> | Volunteer | None | Density | Europe | Native | Native | Relative abundance | na | Habitat was the best predictor of change in faecal density, not hunting pressure. |
| 44 | <i>Vulpes vulpes</i> | Volunteer | Damage | Damage | Europe | Peri urban | Native | Relative abundance | Parasite prevalence | Spotlight shooting did not reduce fox abundance or <i>Echinococcus multilocularis</i> |

| | | | | | | | | | | | |
|----|-----------------------------|---|-------------|-------------|---------------|------------|--------|--------------------|----|-----------|---|
| | | | | | | | | | | | prevalence. Prevalence increased in the area subjected to shooting. |
| 45 | <i>Vulpes vulpes</i> | Volunteer | Density | Density | Europe | Mixed | Native | Density | na | | Effects of increased control effort were largely negated by compensatory immigration. |
| 46 | <i>Vulpes vulpes</i> | Commercial wildlife management contractor | None | Density | Australasia | Mixed | Exotic | Relative abundance | na | | Shooting produced short term reductions and a slow decline over five years. Participants were mostly happy with results. |
| 47 | <i>Vulpes vulpes</i> | Volunteer | None | Density | Australasia | Mixed | Exotic | Density | na | | Shooting mortality was probably offset by rapid immigration. |
| 48 | <i>Vulpes vulpes</i> | Volunteer | None | Density | Europe | Mixed | Native | Relative abundance | | Predation | Spotlight shooting reduced fox relative abundance at two sites in a crossover trial by 45 and 95%. |
| 49 | <i>Felis catus</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Presence | na | | Eradication of cats from a small, tropical island was successful. Shooting was the most effective and efficient removal method. |
| 50 | <i>Felis catus</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Abundance | na | | Intensive shooting began after a major knockdown caused by trapping. Use of detection dogs was critical for locating the last cats and for verifying eradication. |
| 51 | <i>Felis catus</i> | Commercial wildlife management contractor | Eradication | Eradication | North America | Native | Exotic | Presence | | Predation | Poison baiting was the main method used, but shooting and trapping helped remove surviving cats. Cat eradication resulted in decreased sooty tern mortality. |
| 52 | <i>Lynx lynx</i> | Recreational hunter | None | Density | Europe | Mixed | Native | Abundance | | Predation | Recreational hunting can lead to a small reduction in lamb predation when it is sufficient to reduce lynx population size. However, the benefit was very small. |
| 53 | <i>Macropus fuliginosus</i> | Commercial harvest | None | Density | Australasia | Peri urban | Native | Density | na | | Estimated 88% density reduction in an insular kangaroo population. |
| 54 | <i>Macropus rufogriseus</i> | Government | Growth | Growth | Australasia | Native | Exotic | Abundance | na | | Small scale, intensive hunting with dogs was able to reduce local population densities below the 70% level needed for ground shooting to be the most cost effective option. |
| 55 | <i>Macropus rufogriseus</i> | Commercial wildlife management | None | Density | Australasia | Mixed | Native | Relative abundance | na | | Shooting reduced macropod activity on agricultural land, with a diminishing effect with distance from the shooting |

| | | | | | | | | | | |
|----|-------------------------------|---|-------------|-------------|-------------|--------|--------|--------------------|-------------------|--|
| | | contractor | | | | | | | | operation. A companion paper noted that survivors increased their use of agricultural land. |
| 55 | <i>Thylogale billardierii</i> | Commercial wildlife management contractor | None | Density | Australasia | Mixed | Native | Relative abundance | na | Shooting reduced macropod activity on agricultural land, with a diminishing effect with distance from the shooting operation. A companion paper noted that survivors increased their use of agricultural land. |
| 56 | <i>Oryctolagus cuniculus</i> | Government | Eradication | Eradication | Australasia | Native | Exotic | Presence | Native vegetation | Hunting, in combination with trapping, was used after poison baiting and biological control to remove the last rabbits from the island. |

References

- Brennan, M., Moller, H. and Parkes, J. P. (1993) Indexes of density of feral goats in a grassland forest habitat, Marlborough, New Zealand. *New Zealand Journal of Ecology* **17**(2), 103-106.
- Forsyth, D. M., Hone, J., Parkes, J. P., Reid, G. H. and Stronge, D. (2003) Feral goat control in Egmont National Park, New Zealand, and the implications for eradication. *Wildlife Research* **30**(5), 437-450.
- Masters, P., Markopoulos, N., Florance, B. and Southgate, R. (2018) The eradication of fallow deer (*Dama dama*) and feral goats (*Capra hircus*) from Kangaroo Island, South Australia. *Australasian Journal of Environmental Management* **25**(1), 86-98.
- Parkes, J. P. (1990) Feral goat control in New Zealand. *Biological Conservation* **54**(4), 335-348.
- Forsyth, D. M. (1999) Long-term harvesting and male migration in a New Zealand population of Himalayan tahr *Hemitragus jemlahicus*. *Journal of Applied Ecology* **36**(3), 351-362.
- Banko, P. C., Hess, S. C., Scowcroft, P. G., Farmer, C., Jacobi, J. D., Stephens, R. M., Camp, R. J., Leonard Jr, D. L., Brinck, K. W. and Juvik, J. (2014) Evaluating the long-term management of introduced ungulates to protect the palila, an endangered bird, and its critical habitat in subalpine forest of Mauna Kea, Hawai'i. *Arctic, Antarctic, and Alpine Research* **46**(4), 871-889.
- Hothorn, T. and Müller, J. (2010) Large-scale reduction of ungulate browsing by managed sport hunting. *Forest Ecology and Management* **260**(9), 1416-1423.
- Wäber, K., Spencer, J. and Dolman, P. M. (2013) Achieving landscape-scale deer management for biodiversity conservation: The need to consider sources and sinks. *Journal of Wildlife Management* **77**(4), 726-736.

9. Crouchley, D., Nugent, G. and Edge, K. (2011) Removal of red deer (*Cervus elaphus*) from Anchor and Secretary Islands, Fiordland, New Zealand. *Island Invasives: Eradication and Management*. (Eds CR Veitch, MN Clout and DR Towns.) pp, 422-425.
10. Forsyth, D. M., Ramsey, D. S., Veltman, C. J., Allen, R. B., Allen, W. J., Barker, R. J., Jacobson, C. L., Nicol, S. J., Richardson, S. J. and Todd, C. R. (2013) When deer must die: large uncertainty surrounds changes in deer abundance achieved by helicopter-and ground-based hunting in New Zealand forests. *Wildlife Research* **40**(6), 447-458.
11. Fraser, W. (1996) The effect of recreational hunters on deer populations in Pureora Conservation Park. Department of Conservation, Wellington.
12. Fraser, K. W. and Speedy, C. J. (1997) Hunting pressure, deer populations, and vegetation impacts in the Kaimanawa Recreational Hunting Area. Department of Conservation, Wellington.
13. Hagen, R., Haydn, A. and Suchant, R. (2018) Estimating red deer (*Cervus elaphus*) population size in the Southern Black Forest: the role of hunting in population control. *European Journal of Wildlife Research* **64**(4), 42.
14. Husheer, S. W. and Robertson, A. W. (2005) High-intensity deer culling increases growth of mountain beech seedlings in New Zealand. *Wildlife Research* **32**(4), 273-280.
15. Fraser, K. W. (2005) 'Wild deer in Northland: modelling potential new populations and the extant Russell population.' (Landcare Research: Lincoln.)
16. Kaji, K., Saitoh, T., Uno, H., Matsuda, H. and Yamamura, K. (2010) Adaptive management of sika deer populations in Hokkaido, Japan: theory and practice. *Population Ecology* **52**(3), 373-387.
17. Takeshita, K., Tanikawa, K. and Kaji, K. (2017) Applicability of a Bayesian state-space model for evaluating the effects of localized culling on subsequent density changes: sika deer as a case study. *European Journal of Wildlife Research* **63**(4), 63-71.
18. Bennett, A., Haydon, S., Stevens, M. and Coulson, G. (2015) Culling reduces fecal pellet deposition by introduced sambar (*Rusa unicolor*) in a protected water catchment. *Wildlife Society Bulletin* **39**(2), 268-275.
19. Fraser, K. W., Parkes, J. P. and Thomson, C. (2003) 'Management of new deer populations in Northland and Taranaki.' (Department of Conservation: Wellington.)
20. Nugent, G. (1988) Successful control of fallow deer by recreational hunters in the Blue Mountains, Otago. *New Zealand Journal of Forestry Science* **18**(3), 239-252.
21. Martin, J.-L. and Baltzinger, C. (2002) Interaction among deer browsing, hunting, and tree regeneration. *Canadian Journal of Forest Research* **32**(7), 1254-1264.
22. DeNicola, A. J. and Williams, S. C. (2008) Sharpshooting suburban white-tailed deer reduces deer-vehicle collisions. *Human-Wildlife Conflicts* **2**(1), 28-33.

23. Doerr, M. L., McAninch, J. B. and Wiggers, E. P. (2001) Comparison of 4 methods to reduce white-tailed deer abundance in an urban community. *Wildlife Society Bulletin* **29**(4), 1105-1113.
24. Frost, H. C., Storm, G. L., Batcheller, M. J. and Lovallo, M. J. (1997) White-tailed deer management at Gettysburg National Military Park and Eisenhower National Historic Site. *Wildlife Society Bulletin* **25**(2), 462-469.
25. Hygnstrom, S. E., Garabrandt, G. W. and Vercauteren, K. C. (2011) Fifteen years of urban deer management: The Fontenelle Forest experience. *Wildlife Society Bulletin* **35**(3), 126-136.
26. McDonald, J. E., Clark, D. E. and Woytek, W. A. (2007) Reduction and maintenance of a white-tailed deer herd in central Massachusetts. *Journal of Wildlife Management* **71**(5), 1585-1593.
27. Simard, M. A., Dussault, C., Huot, J. and Côté, S. D. (2013) Is hunting an effective tool to control overabundant deer? A test using an experimental approach. *Journal of Wildlife Management* **77**(2), 254-269.
28. Williams, S. C., Denicola, A. J., Almendinger, T. and Maddock, J. (2013) Evaluation of organized hunting as a management technique for overabundant white-tailed deer in suburban landscapes. *Wildlife Society Bulletin* **37**(1), 137-145.
29. Barron, M. C., Anderson, D. P., Parkes, J. P. and Ohukani'ohi'a Gon III, S. M. (2011) Evaluation of feral pig control in Hawaiian protected areas using Bayesian catch-effort models. *New Zealand Journal of Ecology* **35**(2), 182-188.
30. Caley, P. and Ottley, B. (1995) The effectiveness of hunting dogs for removing feral pigs (*Sus scrofa*). *Wildlife Research* **22**, 147-154.
31. Engeman, R., Hershberger, T., Orzell, S., Felix, R., Killian, G., Woolard, J., Cornman, J., Romano, D., Huddleston, C., Zimmerman, P., Barre, C., Tillman, E. and Avery, M. (2014) Impacts from control operations on a recreationally hunted feral swine population at a large military installation in Florida. *Environmental Science and Pollution Research* **21**(12), 7689-7697.
32. Geisser, H. and Reyer, H.-U. (2004) Efficacy of hunting, feeding, and fencing to reduce crop damage by wild boars. *Journal of Wildlife Management* **68**(4), 939-946.
33. Gentle, M. and Pople, A. (2013) Effectiveness of commercial harvesting in controlling feral-pig populations. *Wildlife Research* **40**, 459-469.
34. Gürtler, R. E., Martín Izquierdo, V., Gil, G., Cavicchia, M. and Maranta, A. (2017) Coping with wild boar in a conservation area: impacts of a 10-year management control program in north-eastern Argentina. *Biological Invasions* **19**(1), 11-24.
35. Hanson, L. B., Mitchell, M. S., Grand, J. B., Jolley, D. B., Sparklin, B. D. and Ditchkoff, S. S. (2009) Effect of experimental manipulation on survival and recruitment of feral pigs. *Wildlife Research* **36**(3), 185-191.
36. Keuling, O., Lauterbach, K., Stier, N. and Roth, M. (2010) Hunter feedback of individually marked wild boar *Sus scrofa* L.: dispersal and efficiency of hunting in northeastern Germany. *European Journal of Wildlife Research* **56**(2), 159-167.

37. Krull, C. R., Stanley, M. C., Burns, B. R., Choquenot, D. and Etherington, T. R. (2016) Reducing wildlife damage with cost-effective management programmes. *PLoS One* **11**(1) e0146765.
38. McIlroy, J. C. and Saillard, R. J. (1989) The effect of hunting with dogs on the numbers and movements of feral pigs, *Sus scrofa*, and the subsequent success of poisoning exercises in Namadgi National Park, ACT. *Australian Wildlife Research* **16**, 353-63.
39. Parkes, J. P., Ramsey, D. S. L., Macdonald, N., Walker, K., McKnight, S., Cohen, B. S. and Morrison, S. A. (2010) Rapid eradication of feral pigs (*Sus scrofa*) from Santa Cruz Island, California. *Biological Conservation* **143**, 634-641.
40. Quirós-Fernández, F., Marcos, J., Acevedo, P. and Gortázar, C. (2017) Hunters serving the ecosystem: the contribution of recreational hunting to wild boar population control. *European Journal of Wildlife Research* **63**(3), 57.
41. Servanty, S., Gaillard, J. M., Ronchi, F., Focardi, S., Baubet, E. and Gimenez, O. (2011) Influence of harvesting pressure on demographic tactics: implications for wildlife management. *Journal of Applied Ecology* **48**(4), 835-843.
42. Toïgo, C., Servanty, S., Gaillard, J.-M., Brandt, S. and Baubet, E. (2008) Disentangling natural from hunting mortality in an intensively hunted wild boar population. *Journal of Wildlife Management* **72**(7), 1532-1539.
43. Baker, P. J. and Harris, S. (2006) Does culling reduce fox (*Vulpes vulpes*) density in commercial forests in Wales, UK? *European Journal of Wildlife Research* **52**(2), 99-108.
44. Comte, S., Umhang, G., Raton, V., Raoul, F., Giraudoux, P., Combes, B. and Boué, F. (2017) *Echinococcus multilocularis* management by fox culling: an inappropriate paradigm. *Preventive Veterinary Medicine* **147**, 178-185.
45. Lieury, N., Ruelle, S., Devillard, S., Albaret, M., Drouyer, F., Baudoux, B. and Millon, A. (2015) Compensatory immigration challenges predator control: an experimental evidence-based approach improves management. *Journal of Wildlife Management* **79**(3), 425-434.
46. McLeod, L. J., Saunders, G. and Miners, A. (2011) Can shooting be an effective management tool for foxes? Preliminary insights from a management program. *Ecological Management & Restoration* **12**(3), 224-226.
47. Newsome, T., Crowther, M. and Dickman, C. (2014) Rapid recolonisation by the European red fox: how effective are uncoordinated and isolated control programs? *European Journal of Wildlife Research* **60**(5), 749-757.
48. Tapper, S. C., Potts, G. R. and Brockless, M. H. (1996) The effect of an experimental reduction in predation pressure on the breeding success and population density of grey partridges *Perdix perdix*. *Journal of Applied Ecology* **33**(5), 965-978.
49. Domm, S. and Messersmith, J. (1990) Feral cat eradication on a Barrier Reef island, Australia. *Atoll Research Bulletin* **338**, 1-4.
50. Robinson, S. A. and Copson, G. R. (2014) Eradication of cats (*Felis catus*) from subantarctic Macquarie Island. *Ecological Management & Restoration* **15**(1), 34-40.

51. Rodríguez, C., Torres, R. and Drummond, H. (2006) Eradicating introduced mammals from a forested tropical island. *Biological Conservation* **130**(1), 98-105.
52. Herfindal, I., Linnell, J. D. C., Moa, P. F., Odden, J., Austmo, L. B. and Andersen, R. (2005) Does recreational hunting of lynx reduce depredation losses of domestic sheep? *Journal of Wildlife Management* **69**(3), 1034-1042.
53. Mawson, P. R., Hampton, J. O. and Dooley, B. (2016) Subsidized commercial harvesting for cost-effective wildlife management in urban areas: A case study with kangaroo sharpshooting. *Wildlife Society Bulletin* **40**(2), 251-260.
54. Warburton, B. and Frampton, C. (1991) Bennett's Wallaby Control in South Canterbury. Forest Research Institute, Christchurch.
55. Wiggins, N. L. and Bowman, D. M. J. S. (2011) Macropod habitat use and response to management interventions in an agricultural–forest mosaic in north-eastern Tasmania as inferred by scat surveys. *Wildlife Research* **38**(2), 103-113.
56. Springer, K. (2016) Methodology and challenges of a complex multi-species eradication in the sub- Antarctic and immediate effects of invasive species removal. *New Zealand Journal of Ecology* **40**(2), 273-278.