

# Pasture management for the Burnett catchment



Queensland Government

# Pasture management for the Burnett catchment

compiled by Damien O'Sullivan

First published 2009

Second edition 2012

Third edition 2014

Fourth edition 2022

This publication has been supported by the Queensland Department of Agriculture & Fisheries and Healthy Land & Water, through funding from the Queensland Government's 'Natural Resources Investment Program'. The original publication (2009) was supported through funding from Australian Government's 'Caring for Our Country program'.

© The State of Queensland, Department of Agriculture and Fisheries, 2022. Except as permitted by the Copyright Act 1968, no part of this work may in any form or by any electronic, mechanical, photocopying, recording, or any other means be reproduced, stored in a retrieval system or be broadcast or transmitted without the prior written permission of the State of Queensland, Department of Agriculture and Fisheries. The information contained herein is subject to change without notice. The copyright owner shall not be liable for technical or other errors or omissions contained herein. The reader/user accepts all risks and responsibility for losses, damages, costs and other consequences resulting directly or indirectly from using this information. Enquiries about reproduction, including downloading or printing the web version should be addressed to [DAFCopyright@daf.qld.gov.au](mailto:DAFCopyright@daf.qld.gov.au) (telephone +61 7 3404 6999).

This publication has been compiled by Damien O'Sullivan of the Drought Policy and Response group, Department of Agriculture and Fisheries. The author acknowledges contributions from Ian Crosthwaite, Bill Edwards, Col Paton (EcoRich Grazing), Gavin Peck (DAF), Louise Walker (DAF), Giselle Whish (DAF), Jodie Ward (DAF), Greg Bath (DAF) and Bruce Lord (Healthy Land and Water).



Introduction .....	1
<i>Key pasture management skills</i>	
Pasture basics .....	2
<i>Managing pasture condition</i>	
Grazing systems .....	8
<i>Grazing to suit your situation</i>	
Enhancing native pastures .....	11
<i>Using legumes to increase carrying capacity</i>	
Establishing sown pastures .....	14
<i>Best option for old cultivation land</i>	
Burnett land types.....	20
<i>Selecting the best pasture mix</i>	
Legumes for Burnett pastures.....	53
<i>Descriptions and planting rates</i>	
Grasses for Burnett pastures.....	65
<i>Descriptions and planting rates</i>	
Further information.....	78
<i>Contacts and other information sources</i>	
Workshops .....	80
Species list.....	81



# Introduction

The beef industry is a major contributor to the wealth of the Burnett region and this valuable industry relies on productive pastures. This booklet aims to help land managers better understand the development and management of pastures suited to the region.

Selecting the best mix of grasses and legumes for each land type will ensure the best financial return on investment from improved pastures. This booklet is designed to help all grazing property managers, from those on small grazing blocks to properties of 10 000 ha or more.

The Burnett catchment lies within the southern black speargrass region of Queensland's pasture communities. On the surface it may seem simple to manage a paddock full of grass but the way graziers manage stock numbers under a variety of climatic conditions can dramatically improve or deteriorate the condition of the ecological system.

To establish and maintain healthy and sustainable pastures, graziers need to know:

- what plants are in the pasture and the condition of the pasture
- whether there are changes occurring in the numbers of broadleaf weeds, woody weeds or other unfavourable species
- how many stock to carry with the feed on hand (i.e. when is it time to sell stock or begin to feed supplements?)
- how changes in stock or pasture management will affect the pasture composition or condition
- how to maximise income without jeopardising pasture health
- how to demonstrate that the property management strategies are sustainable
- how to use a system like *Stocktake* to forage budget and monitor changes in pasture condition over time
- manage sown pastures to minimise spread into ecological sensitive areas.



# Pasture basics

Pastures are complex ecosystems. A wide variety of ecosystem processes and management decisions affect the composition and productivity of a pasture over time.

## Pasture composition

The composition of pastures varies considerably. It is important for managers to have a clear idea of what they expect from their pastures and choose plant species that have the capacity to fulfil grazing requirements.

There are two very broad plant types found in most pastures:

- annuals and weak perennials—plants that live for one or a limited number of growing seasons
- perennials—plants with variable life spans, from several years to decades.

In the grazing situation the main plants of interest are:

- grasses—low-growing vegetative plants that can tolerate grazing. Stoloniferous grasses like couch and Rhodes grass spread by seed and above-ground roots called stolons. Grasses like kikuyu have stolons and underground roots called rhizomes. Tussock grasses grow in clumps as distinct individual plants
- forbs—non-woody broadleaf plants

- legumes—plants with a varying ability to fix nitrogen with the help of bacteria in root nodules (e.g. clovers)
- sedges—grass-like plants (e.g. nutgrass)
- woody weeds—woody plants with the ability to compete with pasture plants for sunlight, nutrients and moisture (e.g. wattles)
- other weeds—any undesirable plant in a pasture that affects carrying capacity of the pasture due to its unpalatability, competition with more favourable species or toxicity to livestock (e.g. lantana and African lovegrass).

## Features of a healthy pasture

A healthy pasture has a:

- high percentage of palatable, productive, perennial grasses (3P grasses)
- productive legumes
- small number of annual plants (e.g. small burrgrass)
- small number of weeds (e.g. verbenas)
- high frequency of desirable forbs (e.g. native legumes such as glycine)
- variety of other favourable species.

## What affects pasture condition and quality?

Climate, soil type, plant type and grazing pressure all interact to determine the quality, productivity and sustainability of pastures.

## Climate

Climate, and rainfall in particular, has a major effect on pastures. Given the wide variation in climatic conditions between years, it is unreasonable to expect pasture bulk and quality to remain constant.

The nutritive value of a plant varies with seasonal conditions and soil fertility. In years of high rainfall, plant yield is often high but protein content is low. The opposite is often the case in lower rainfall years. Frost, wind, humidity and heat can also affect pasture quantity and quality.

The effectiveness of rainfall is the major determining factor for pasture growth. A combination of the following factors will determine the amount of effective rainfall and subsequent pasture growth:

- ground cover—pasture condition, plant type and species (see Table 1)
- soil type and condition—aggregate structure, pore size, amount of organic matter, carbon and minerals
- rainfall intensity
- evaporation
- slope
- tree cover.

Table 1 indicates that the perennial tussock grasses grew more dry matter per mm of rainfall.

## Soil type

Generally, the better the soil, the better the pasture. While we often look at a particular grass as being very good for stock feed, the protein level of that grass varies depending on soil fertility. Low soil phosphorus and nitrogen are generally the most common nutrient restrictions on pasture growth.

Organic matter is an important soil component, providing plant nutrients, binding soil aggregates, improving infiltration and feeding soil micro and macroorganisms. Low organic matter levels inhibit plant growth.

Once pasture cover drops below 90% there is a serious risk of soil erosion due to increased run-off.

Much of the fertility in Burnett soils is carried in the topsoil so it is important to manage pastures in a way that will preserve and improve the quality of this soil zone. A paddock with pasture composed of annuals and large areas of bare soil loses significant quantities of fertile topsoil every year (see Figure 1).

Table 1 – The effect of pasture type on water infiltration and dry matter production

	Water infiltration %	kg dry matter/ mm rainfall
<b>Perennial tussock grasses</b>	76	4
<b>Stoloniferous grasses</b>	55	3
<b>Annual grasses</b>	40	2
<b>Bare ground</b>	25	0




Figure 1 – Soil losses from pastures.

### Plant type

Each type of plant community has something different to offer land managers. Three main pasture types contribute to the Burnett catchment.

#### Native pastures

These pastures are often underused and undervalued. They are a cheap resource, costing 3–18c per kg of liveweight gain in cattle compared to around \$3.20 per kg of liveweight gain in cattle on a grain diet.

Native pastures are still the main pastures supporting Queensland's grazing industry. They can benefit from a careful burning regime and, with proper management, they can be productive and profitable.

#### Native pastures + legume

Oversowing legumes into native pastures is a cost-effective way of increasing production. Stylos and Wynn cassia are commonly used in the Burnett. Clovers and medics can perform well in wet winters.

### Sown pastures

Pastures sown into a prepared seedbed are the most productive, and the most expensive. These pastures will usually run down after 5–8 years due to a lack of nitrogen. They can be maintained for longer with fertiliser applications and careful renovation. Legumes are an essential addition to pastures to reduce the effects of run-down.

### Grazing

Stock will select the highest quality diet available to them. Increasing the stocking rate reduces each animal's ability to select and so reduces the quality of the diet. Forcing stock to eat more than leaf (e.g. stem) will reduce the weight gain per head.

About 25–40% of a pasture grass plant (above the ground) is leaf. To maintain strong pasture plants do not allow animals to use any more than a third of the grass growth over 12 months. Using about one-third of the feed on offer gives stock the best quality diet and avoids overgrazing the pasture (see Figure 2).




Figure 2 – Using just one-third of the feed on offer for stock helps maintain quality pastures.

If stock are allowed to continually eat green shoots as they emerge, the plant is unable to build up carbohydrate reserves in its roots. This causes root loss and a loss in plant productivity (see Figure 3).

As the number of stock grazing on a pasture increases, the potential liveweight gain decreases (see Figure 4). Increased stocking pressure results in the loss of more palatable pastures species and a decline in pasture quality.

### Improving pasture palatability and productivity

Several factors influence pasture palatability and productivity. Use the following principles to help manage pastures to achieve long term productivity.

#### Leafiness

Leafiness and the protein content of the leaves determines the palatability and productivity of grasses. Nitrogen content (and so protein) varies




Figure 3 – An over-grazed plant compared to a moderately grazed plant. Note the difference in root mass.




Figure 4 – Weight gain (annual liveweight gain in kg/head vs stocking rate head/ha).

between plant stem and leaf and the time of the year (see Table 2) and leafiness varies with pasture species (see Table 3).

Perennial grasses live for more than a year, providing bulk and cover year round if managed properly. If perennial grass cover declines, the result is inefficient energy capture, loss of nutrients to the plant, inefficient use of rainfall and a decline in soil condition. So with less pasture production per mm of rainfall there is a decrease in stock carrying capacity.

#### Age of pasture plants

Most pastures have maximum energy and protein for a period of 40–60 days. Unless there is continued pasture growth there is an inevitable decline in the value of the pasture. Figure 5 shows the decline in crude protein and digestibility of black speargrass over time following rainfall.

Table 2 – Nitrogen content % of black speargrass leaf and stem

	Early wet season %N/kg dry matter	Mid dry season %N/kg dry matter
Leaf	10.1	4.1
Stem	3.6	1.1

Table 3 – Leaf percentage of the above-ground portion of some grass species

Grass	% leaf
Green panic	30-45
Speargrass	30
Wiregrass	10
Golden beard grass	high leaf % but low yield

### Legumes in the pasture

Legumes provide an extra source of protein for animals grazing the pasture as well as contributing nitrogen to the soil for grasses in the pasture to utilise. Compared to grasses they also retain their digestibility for a longer period of time (see Figure 6).

This is important as a pasture with at least 25% legumes (4 plants/m<sup>2</sup>) will help animals gain weight for a longer period in the dry season.

### Spelling pastures

Grasses need a period of rest if they are to set seed and produce to their maximum potential, especially after burning or heavy grazing. Spell pastures over the grazing season for at least six weeks or until the grass goes to seed.




Figure 5 – Change in crude protein and digestibility over time.




Figure 6 – Digestibility of legume vs grasses.

If you are spelling pastures to set legume seed a longer time period is needed. This may need planning over a number of years to ensure every paddock receives a spell over time.

Lack of rest for pasture over the growing season is the main reason for loss of favourable pasture species.



Perennial stoloniferous grasses provide ground cover, protecting soil from erosion.

## Fire

Fire can be a useful tool for pastures if used correctly. In the speargrass region burning every 3–4 years is recommended. This however will depend on the season, stocking rates and the grazing intention for the pasture. Most of the vegetation in this region has developed under a regime of fire. Grasses like black speargrass are adapted to fire and can increase in density with correct and timely burning.

Carry out burning in spring when ground litter is damp and will not readily burn. Choosing the right time to burn can be difficult and dangerous. Remember a fire permit is required for any burning. Spell burnt pastures until they go to seed. The green pick after burning pasture is very palatable and nutritious to stock but grazing at this time will damage 3P grasses.




Figure 7 – Tree basal area vs pasture yield.

## Timber and woody weeds

Woody weed density affects pasture growth. Generally, the less trees and woody weeds the more water is available for pasture growth (see Figure 7). However in the very long term a 20% canopy cover in the pasture is the optimum for maximum production and sustainability.

## How to set carrying capacity

**Step 1** – Determine the amount of feed available using pasture standards or pasture cuts.

A *Stocktake* workshop will show you how to do this (see page 61).

**Step 2** – Aim to utilise between 25–40% of feed grown to give stock the best quality diet and allow grasses the opportunity to grow to their best potential (how much feed do the stock need? or what are the triggers for removing stock from a paddock?).

**Step 3** – Budget to leave a proportion of the feed to provide for trampling by stock, ground cover, decomposition and for other fauna (Figure 2).



*Fodder conservation can supplement animals during periods of standing feed shortage.*



# Grazing systems

There are many grazing management systems available for stockowners to implement. Each system has advantages and disadvantages. If used incorrectly, any of these grazing systems can cause damage to the pasture and land resource base.

The most common grazing systems are described below. Although the names used for these systems varies considerably within the industry, these descriptions will generally hold true.

## Set stocked / continuous grazing

This is a traditional grazing system where the same number of cattle are kept in a paddock all year, every year. A good example is a breeder paddock where an owner may aim to run 50 breeders each year. In dryer years cows will have to be fed or culled to cope with the shortfall in feed. In better years cow numbers could be increased but in most cases it would be better to allow the extra growth of pasture and enable the more favourable species to go to seed.

With set stocking it is common for some areas of the paddock to be preferentially grazed while areas that are less accessible or further from water may experience very little grazing pressure. If stock numbers are kept at recommended levels for the land and pasture type this can be a low labour input system.

## Wet season spelling

This system involves removing all stock from one or more paddocks (spelling) on a regular basis over the wet season. Research has shown wet season spelling is one of the most effective ways to maintain a healthy and productive pasture. A joint CSIRO/DAF study found that this method of pasture management was a common factor in many successfully managed beef operations studied across Queensland.

The recommendation is to remove stock from a paddock for at least six weeks over the growing season. This time period may vary depending on rainfall, with a longer time period needed in some years. Essentially, the aim is to allow the better 3P (palatable, productive and perennial) grass species to go to seed. Over a period of several years each paddock on the property can be spelled on a rotation.

## Rotational grazing (time based)

Under the timed rotational grazing system cattle are moved to a new paddock after a pre-determined period of time. Generally cattle on the property are grouped together into large mobs and each mob is moved in a rotation around all available paddocks. Paddocks on most properties vary in size so bigger paddocks will be grazed for longer whilst smaller paddocks may only be grazed for a few days. This system allows paddocks to spell and the bigger numbers of stock in a paddock spreads the grazing pressure away

from favoured areas such as creek flats or areas of heavier soil types. An advantage of the rotational system is that stock movements can be timed to pass through the cattle yards for routine animal husbandry jobs. This can reduce the time needed for mustering several mobs held in a number of paddocks, as is the case with set stocking. Provided the water supply in each paddock is adequate for the increased stock density, there is no need for additional fencing or infrastructure.

### Rotational grazing (seasonal)

This system is similar to timed rotational grazing but the cattle are moved from paddock to paddock on a timing that is based on pasture growth and the season. Generally, paddock moves will be more frequent in the wet season in an attempt to keep pasture in a vegetative state rather than letting it go to seed. This can promote plant tillering and help to increase the amount of leaf each plant produces. In winter, when grass growth has finished the cattle move less frequently.

Like time based movements in a rotational system this method allows the spreading of grazing pressure over a larger area of the paddock, reducing patch grazing. This system has the advantage of taking into account times when pasture is susceptible to damage and using spelling to allow grasses to go to seed. During winter when grass growth has essentially stopped the longer grazing periods do not cause damage to the pasture. Provided

the water supply in each paddock is adequate for the increased stock density, there is no need for additional fencing or infrastructure.

### Cell grazing

This system of grazing has a centralised watering point and uses electric fencing creating a set of cells in a wagon wheel or block formation. Under this system a single paddock is cut up into multiple cells and the cattle are moved through the cells every 2–3 days, depending on pasture growth. Like the rotational grazing system, wet season paddock moves will be more frequent in an attempt to keep pasture in a vegetative state rather than letting it go to seed. During winter the time between moves can be increased.

Generally, managers using this system use a grazing chart to record the number of adult equivalents they have grazing over the area and this will give them a number of stock days/ha. By combining this information with rainfall records, the manager can estimate the feed grown per millimetre of rainfall.

This system allows for very accurate estimates of how much feed is available for the coming season and enables the operator to make early decisions about stocking rates (e.g. when to buy or sell).

Grazing charts from previous years are an invaluable tool when making future management decisions. With detailed information about paddock carrying capacity the operator can optimise pasture utilisation. The

system of cells needs to be carefully planned and may need significant investment in single wire electric fencing and water infrastructure.

### High density grazing

This is a very intensive grazing system where cattle are grazed at high stocking rates and moved daily or 2–3 times/day. An example stock density might be 740 animals, each weighing 450 kg, on 1 ha (300 x 450 kg animals on 1 acre) for 24 hours. The stock are moved at least daily with some operators moving the stock up to eight times a day. Electric fencing is used and in many cases water troughs are moved with the stock.

Proponents of this system suggest that stocking rates of 50% more animals are possible, compared to other grazing systems. This system aims to mimic the effects of large grazing animals in Africa and buffalo in the USA. Originally these animals moved quickly as they migrated across an area, eating everything in their path and fertilising the pasture with their manure and urine. These rangeland pastures were then rested for considerable time periods before the herds returned.

High density grazers generally aim for the cattle to consume 60% of the pasture, trample 20% into the soil and leave 20% standing. The trampling of the pasture into the soil gives soil biology the organic matter it needs to increase soil carbon levels. One noted proponent of the system

claims a dramatic increase of soil carbon from 1.5% to 8% over eight years. A key to this system is the long rest periods for the pasture. In more fertile areas an area of pasture may only be grazed for a total of 4 days/year. In drier environments each section of pasture may only be grazed once a year. Graziers have noted a vast improvement in soil health and biology after using the system for three years. Research in Texas USA found that pasture composition could be beneficially changed in four years with a carefully managed high density grazing system.

### In summary

Research in northern Australia has consistently shown that the major opportunities for improved land condition and productivity lie in:

- better spatial distribution of grazing pressure (through location and number of water points, sub-divisional fencing)
- better matching of stocking rate with carrying capacity
- targetted use of wet season spelling.

All grazing systems have their advantages and disadvantages. Each stockowner needs to carefully examine each system and adopt a method that suits their management, labour resources and ability to invest in the necessary infrastructure. Ultimately the aim of any grazing system is to improve the health of the pasture and soil resources.



## Enhancing native pastures

Legumes increase the quality of native pasture and so increase carrying capacity, breeder performance and animal growth rate.

Introducing legumes can benefit native pastures and run-down improved pastures, increasing carrying capacity and property income. Legumes supply some nitrogen to the soil and boost grass growth, but their main benefit is in offering a higher protein diet to the grazing animals. The protein level of summer growing grasses drops as plants mature and remains low during autumn and winter. Legumes, with their high protein content, are an excellent source of nutrition for stock and maintain their levels of protein much longer than grasses (Figure 6).

Most native pastures in the Burnett have one or more native legumes present. Rhizobia on the roots of the legume fix nitrogen and over time this is made available in the soil to be utilised by other plants. This helps to minimise the run-down in production that usually occurs in fertilised sown pastures after 5–6 years.

Many legumes have deep tap roots and are able to access moisture and nutrients deep down in the soil. The Burnett area is generally more suited to tropical and sub-tropical legumes

such as Wynn cassia and stylos. However in wet winters clovers and medics will grow.

## Grazing improved native pastures

South Burnett case studies show that, depending on soil type, graziers can rely on the following 'rules of thumb' when planning to add legumes to their pastures:

- Some legumes will persist. Stylos have been established in some pastures for 20 years or more. At least 25% legume in the pasture sward is desirable.
- Under good summer conditions legumes in a pasture will give an extra 0.1 kg per head per day over the 0.7 kg for grass-only pastures.
- Animals usually lose weight during winter when grazing native or improved grass-only pastures. However in mild winters, legumes such as lotononis, white clover and lucerne in pastures can add an extra 0.5 kg per head per day over the 0.3 kg expected for grass-only pastures.

Manage pastures with legumes carefully as Wynn cassia and some stylos have low palatability and can dominate pastures if grass is grazed out, causing an overall loss of production.

Clovers, medics and lucerne can also cause bloat in cattle when they make up a large proportion of an animal's diet.

Case studies have shown that introducing legumes can increase stocking rates by up to 20%. Of the extra income that legumes generate, 90% comes from the increased stocking rate and only 10% from the extra weight gain per beast. However, long term stability of the pasture should be considered. Many properties are already stocked to capacity even with some degree of pasture improvement.

### Establishing legumes into existing pastures

Legumes can be introduced into existing native or sown pastures in a variety of ways as outlined below. Recommended planting rates vary considerably depending on the planting mix and composition of the existing pasture (see page 54).

All legumes need to be inoculated at planting to ensure the correct species of rhizobia is available to fix



*The addition of legume to a productive native pasture will often provide a worthwhile boost to livestock and pasture alike.*

nitrogen from the atmosphere through nodules that form on the roots. The inoculation process involves mixing the inoculant with a sticker and the seed within 24 hours of planting. The inoculant is a living material and requires careful treatment. Follow the recommendations that come with the inoculant and use it before the expiry date, after which time viability is lost.

Many seed companies inoculate legume seeds as part of the coating process. In this case, fresher coated seed is recommended.

### Renovation

The grass pasture is very roughly cultivated and legume seed sown onto the surface. As the grass regrows it competes with the legume seedlings and losses can be very high. This method is best where temperate legumes, such as clovers and medics, are sown into grass pastures in autumn. Frost will reduce the grass competition. Tropical legumes can be established this way but it is usually less successful.

### Ploughed strips

Legume seed is sown on the surface of ploughed strips. The legume then spreads to the un-ploughed area by seed or runners. Usually the strips require two workings to reduce the grass competition. Strips should be ploughed on the contour and not across gully lines. Up to 20% of the area may be disturbed to introduce the legumes. Depending on the legume chosen, kangaroos, hares and rabbits can cause significant damage to the establishing legume plants.

### Bandseeding

A bandseeder plants, fertilises and sprays herbicide in strips in one pass. Bandseeding is better than preparing a cultivated seedbed because it is cheaper, there is less risk of soil erosion and the pasture is out of production for less time.

Only one third of the total area is sprayed. The herbicide reduces grass competition and so improves seedling survival and growth.

Another system used on various implements is a broad flat foot on a tyne. The foot cuts under the existing pasture, eliminating grass competition for the pasture seeds sown into the ploughed area.

### Sod-seeding or direct-drilling

Conventional sod-seeders do not handle rough country as well as a bandseeders do. They have difficulty in placing small seeds near the surface and do not remove grass competition without a separate spraying operation. Best results have been from sod-seeding temperate legumes (lucerne and medics) in autumn. Many direct drills have the ability to plant into pastures. Larger seeded legumes such as siratro are more likely to be successful.

### Crocodile seeder

This is an implement with a large cylindrical drum to which shovel-like tools are welded. Seed placed in the drum escapes through holes at each shovel as the implement is towed along. It is a cheap method of establishment and a large 4WD can tow the implement.

### Surface sowing

Seed spread on the surface without preparing a seedbed is often wasted. It has to compete with the established pasture and removal by ants. The best chance of success is using medics and clovers in autumn when frosts kill off the competition and good rain is expected.

There are some good examples of lotononis sown by air at very low rates. These pastures have taken seven to ten years to become highly productive. Surface sowing is an option for seeding inaccessible areas. For example, seeds may be broadcast from the air or by hand into hilly country after a fire. Success depends on good rain after planting.

### Animals

Seed fed to cattle can be spread in the manure. This is usually not economical given current seed prices and losses of highly digestible seeds.

A trial showed that there was more reliable seed strike around the location of movable feed troughs than in manure.

Alternatively a small plot of pure legume can be grown, grazed when seed is mature then the animals released into a suitable paddock. Hundreds of dollars of seed can be 'harvested' in this way. This is a very effective way of spreading hard seeded tropical legumes such as the stylos (e.g. fine stem stylo).



## Establishing sown pastures

Sown pastures have the potential to improve stock production in many areas of the Burnett. Introduced grasses and legumes can rejuvenate run-down native pastures and old cultivation country.

It is not recommended to replace productive native pastures of black speargrass, Queensland blue grass or forest blue grass unless they have been overtaken by unpalatable species such as wire grass. Careful management of native pasture provides the cheapest long-term fodder source for stock. The addition of a tropical legume may be all that is needed to give native pasture a worthwhile boost (see ‘Enhancing native pastures’ on page 11).

New sown pastures will be the most productive for the first 3–5 years. After this, there is usually a run-down of nitrogen in the soil and production decreases. To maintain production levels, use fertiliser, mechanical pasture renovation and/or the addition of legumes.

The rules of thumb for establishing sown improved pastures are:

- the better the seedbed, the better the establishment
- sow most grass and small legume seeds on the surface
- sow when there is good soil moisture at depth
- know the quality of the seed before planting.

## Selecting pasture mix

The mix of grasses and legumes planted depends on the soil or land type and on the cost and availability of seed. The ‘Legumes for the Burnett’ and ‘Grasses for the Burnett’ sections (pages 53 and 65) provide detailed information to help select suitable species. Preparation time and planning is important, it is best to plan up to 12 months in advance for a pasture planting program.

## How much seed to plant

Aim to plant 2–3 kg/ha total of grass seed and 2–3 kg/ha total of legume seed. The amount of seed you decide to plant will depend on the cost and availability of seed. Obviously the more seed you can afford to plant the better the establishment is likely to be. Keeping grazing animals off the newly planted pasture until it goes to seed will mean much more seed will be added to the seed bank.

## Pure live seed (PLS)

All commercial seed should be sold with a ‘statement of seed analysis’ label showing the independent laboratory analysis of the seed, including the germination percentage and a seed purity percentage. The germination percentage indicates the amount of fertile pasture seed in each kilogram you purchase. The purity percentage indicates how much inert matter or other seeds (e.g. weeds) are present in that batch of seed. Using this information you can determine the ‘pure live seed’ percentage (PLS %) of the batch of seed you are purchasing. This will help you choose

the best value seed available and work out the required planting rate. Aim to sow at least 1 kg of PLS/ha. To determine how much pure live seed you are getting, use the formula:

$$\frac{\text{Germination \%} \times \text{Purity \%}}{100} = \text{PLS \%}$$

In some cases the cheapest seed may not be as cheap as it seems. For example,

Grass seed sample A costs \$15/kg. Its germination percentage is 20% and the purity percentage is 80%.

$$\frac{20\% \times 80\%}{100} = 16\% \text{ PLS}$$

$$\begin{aligned} \text{Real cost} &= \$15 \times 100/16 \\ &= \$93.75/\text{kg PLS} \end{aligned}$$

Grass seed sample B costs \$18/kg. Its germination percentage is 40% and the purity percentage is 90%.

$$\frac{40\% \times 90\%}{100} = 36\% \text{ PLS}$$

$$\begin{aligned} \text{Real cost} &= \$18 \times 100/36 \\ &= \$50/\text{kg PLS} \end{aligned}$$

### Coated seed vs uncoated seed

Fluffy seeds such as Rhodes grass and creeping bluegrass can be problematic to sow using commonly-used equipment. This problem can be overcome by using coated seed. Generally you will need to plant more coated seed per hectare than if the seed was uncoated because the coating forms part of the weight. Ask your seed merchant for the recommended planting rate that allows for the coating when buying coated seed. Seed coatings can vary

widely in the ratio of seed to coating so it can be difficult to compare different sources of seed. Many companies now have proprietary seed coating mixtures with additives to help seedling establishment. Insecticides can also be added if ants are collecting and burying seed. In the case of legumes the coating can also include the rhizobial inoculum needed for the legume to fix nitrogen, saving time. Some Independent trials have shown better germination rates with uncoated seed.

### Planting times

When to plant is the most difficult decision in establishing pastures. It is always hot and dry after you plant!

Plant summer pastures from August to March. Early or late is the best option. Early rain in September will be very useful for germination and establishment, while rain may fall more reliably in January–February. Hot, dry spells from October to December often kill young seedlings.

April–May is the preferred time for temperate species such as lucerne, medics, clovers and vetches.

Grass weeds, particularly in old cropping ground, can cause establishment failures. Sometimes the pasture will not be at its best until the second year. Weeds are generally worse in spring plantings. Delaying planting until January–February is often worthwhile but do not leave it so late that short new pasture is frosted. In a grass/legume pasture mix, broadleaf weeds can compete

with the planted pasture. In some cases slashing after good pasture root development can help minimise broadleaf weeds.

Pasture seeds are usually planted dry and will be dormant until the next rain. The seed zone needs to be moist for three to four days for germination to occur. Along with surface moisture, pasture establishment relies on deeper, subsoil moisture.

## Cover crops

Cover crops have been widely used to establish pastures in the Burnett. Cover cropping involves planting crops such as maize, oats, grain sorghum, forage sorghum and millets at half their normal seeding rates.

There are four reasons given to justify the use cover crops:

- shading to protect young seedlings on sandy soils
- suppressing weeds
- providing grazing or some cash return if the pasture fails
- reducing erosion on steep country.

Of these reasons, only minimising erosion is valid. Planting in early spring or autumn will avoid seedlings being burnt in sandy soils. Planting a cover crop thick enough to suppress weeds also suppresses the pasture.

A cover crop is often grazed before the pasture has fully established. Successful pasture establishment using cover crops occurs only in good seasons. Often the cover crop will compete for moisture with the pasture, inhibiting its growth and establishment.

Cover crops can be used as an intermediate crop to reduce weed burden when a permanent pasture is planted. A large variety of annual seeds can be planted to help improve soil condition.

## Seedbed preparation

Seedlings need some soil disturbance to establish. Few legumes and almost no grasses will establish in undisturbed soils, especially those with hard-setting surfaces.

Failure of sown pastures is often due to poor seedbed preparation so allow ample time to prepare a suitable seedbed. If preparation is left too late the result is often a rough, cloddy seedbed, poor weed control and little subsoil moisture. Prepare a firm seedbed for small pasture seeds.

Avoid over-cultivation of soils that are prone to setting hard or crusting after rain. These soils include many Burnett forest soils and some old cultivation soils that are often poorly structured and have a tendency to surface seal due to lack of organic matter.

Seedbed preparation does leave soils prone to erosion so try to follow these guidelines to reduce soil loss:

- do not cultivate in gullies and drainage lines
- divert run-off water away from the cultivated ground
- leave grass strips in ploughed areas

- avoid over-working soils to produce a very fine, powdery seedbed
- if possible cultivate on level contours in the paddock.

### Planting rates

Planting rates for legumes and grasses are provided on pages 54 and 66 respectively.

### Planting methods

The planting method chosen depends on the seed type and machinery available. Often a neighbour with good pastures has the best experience for your local area. Seed can be spread using a rolling drum seeder, fertiliser spinner or combine drill with the seed hoses removed. Other options are full cultivation, direct drill, sod seeding, bandseeding and crocodile seeder.

Adding a legume during pasture cropping can also have some success (see ‘Enhancing native pastures’ on page 11).

#### **Planting fluffy seeds**

Fluffy seeds, such as Rhodes grass and creeping blue grass, are hard to spread with most planters. To make planting easier try mixing the seed with fertiliser, cracked grain or sawdust to act as a carrier.

Another option is to purchase coated seed. If mixing with fertiliser, do not leave seed and fertiliser mixed longer than necessary. If the planting mix includes inoculated legume seed, these seeds must be pelleted before mixing with fertiliser.



*Fluffy seeds, like creeping bluegrass, are easier to plant if the seed is coated.*



*Roller or drum seeders are necessary to sow uncoated, fluffy grass seeds such as buffel, Rhodes grass and creeping blue grass.*

Mixing seed with sieved, dry sawdust is safer than using fertiliser as the carrier. Use twice the volume of sawdust to grass seed. Calibrate the seeder using only sawdust, before adjusting with seed and sawdust. If broadcasting, only fill the hopper with enough seed for 1–2 ha to prevent bridging. Use runs 1.25–1.5 m apart and check how far the seed has thrown. It may be better to plant across the wind.

Most seed companies now offer only coated seed. Using coated seed



*Plant pangola grass as runners because the seed is generally infertile.*

will generally make planting easier and more accurate. The coatings are lime based and can contain various fertilisers. Remember to increase the planting rate to compensate for the increased weight and volume of the coated seed.

Coated seed is advantageous if ants are likely to remove or destroy fluffy seeds. The seed coating can have insecticide added to protect the seeds from ant damage.

In the case of legumes, the rhizobia inoculant can be added when the seed is coated and will remain viable for 28 days.

### Planting depth

Seed is usually placed on the surface and lightly covered with soil. To achieve the best strike with most pasture grasses and legumes do not bury the seed more than 10 mm deep. However, lucerne, purple pigeon grass and silk sorghum can be planted as deep as 25 mm. Many pasture establishment problems are caused by planting too deep.

A small seeds box or a C-seeder mounted on a combine will allow planting at two depths.

### Rolling and harrowing

Rolling and/or harrowing will improve emergence on most soils. However, many old cultivation soils are poorly structured and tend to surface seal. Using a roller after planting can make crusting worse.

Harrows with tynes upright or chains and weldmesh will cover seed from 0 to 10 mm deep, depending how rough the seedbed is. Very loose soils, such as red snuffy soils, may need rolling before and after planting. Rolling with dual tractor wheels has proven very successful on these red soils.

### Fertiliser

Old cultivation soils often have low fertility. Before investing in pasture improvement it is worth getting a soil test to determine if there are any problems. A fertiliser program can then be implemented. Phosphorus, nitrogen and potassium are the main nutrients pastures need. Many red soils in the Burnett are very acidic. These soils



*Well managed buffel grass and seca stylo pasture.*

will benefit from applications of lime before planting the improved pasture. Lack of nitrogen is a common cause of poor grass growth and low nutritive value, especially on some old, cultivated forest soils.

## Weeds

Weeds can be a major problem when establishing pastures. They are usually worse in old cultivation ground.

Ways to reduce competition from weeds include:

- Delay planting until January–February to reduce the number of grass weeds.
- Slash to reduce weeds, this will limit their growth and reduce competition for moisture.
- Use herbicides to control grass weeds. Most herbicides will damage young legumes.



*Healthy, recently established Rhodes grass pasture.*

- If broadleaf weeds are a major problem, another option is to plant grasses and then when they are established oversow with legumes. This way broadleaf herbicides can be used in the establishment phase of the grass.

## First year management

Grazing management in the first year is critical to the establishment and long-term viability of sown pastures. There are two general approaches:

1. no grazing until the pasture seeds
2. lightly grazing once or twice during the first summer–autumn.

Light grazing is preferable. The young pasture will provide some feed, and grazing will encourage the grasses to spread out and reduce the build-up of excessive growth that can smother the next generation of self-sown grasses and legumes. Heavy grazing will permanently damage the pasture.

Allowing the pasture to seed in the first year is important only for grasses that reproduce only via seed rather than runners. Rhodes grass and creeping blue grass will fill in the gaps without seeding.

Frosted pasture can be grazed. After spring rain, allow a build-up of feed before grazing. Planning a forage sorghum crop for this time can give pastures a break.

More pastures are destroyed in the first year due to overgrazing than for most other reasons. Sown pastures are not indestructible.



# Burnett land types

There are 32 land types identified for the Burnett area. Descriptions of characteristic land resource areas, landforms, woody vegetation and expected native pasture species for each main land type are provided in this section.

## Inland landtypes

### Alluvial landtypes

Blue gums on cracking clay.....	22
Blue gums on loam & duplex.....	24
Box on clay.....	25

### Scrub landtypes

Bastard scrub.....	21
Softwood scrub.....	38

### Open forest landtypes

Blue gums on granite.....	23
Box on erosive soil.....	26
Brigalow & brigalow belah.....	27
Brigalow with melonholes.....	28
Gum-topped box.....	29
Ironbark & bloodwood on non-cracking clay.....	30
Ironbark & spotted gum on duplex and loam.....	31
Ironbark on basalt upper slopes and benches.....	32
Mixed open forest on duplex and loam.....	33
Narrow-leaved ironbark on granite.....	34
Narrow-leaved ironbark & wattles.....	35
Silver-leaved ironbark on cracking clay.....	36
Silver-leaved ironbark on granite.....	37
Spotted gum ridges.....	39
Tall open forest on snuffy soils.....	40

## Coastal landtypes

### Alluvial landtypes

Bloodwood & stringybark.....	41
Blue gum flats.....	42
Tea tree flats.....	52

### Scrub landtypes

Hoop pine scrub.....	45
Softwood scrub.....	51

### Open forest landtypes

Blue gum, ironbark & bloodwood slopes and hollows.....	43
Gum-topped box.....	44
Ironbark, stringybark & supplejack ridges.....	46
Ironbark & bloodwood on non-cracking clay.....	47
Ironbark & blue gum on basalt ridges.....	48
Ironbark & spotted gum on duplex and loam.....	49
Mixed eucalypts on uplifted coastal plains.....	50

## Bastard scrub



<b>Land resource area</b>	Volcanic uplands, Red tablelands.
<b>Landform</b>	Ridge crests; broad ridges; some scarp areas; upper, mid and lower slopes of undulating rises and low hills. Commonly slopes 3–10%, occasionally steep 10–25% slopes, and in minor areas as steep as 45%.
<b>Woody vegetation</b>	Open forest to closed scrubs of softwood species (vines, bottle trees, white cedar, crow’s ash, figs) and / or hoop pine and / or narrow-leaved ironbark open woodland. Other species may occur include Burdekin plum, Yarraman ironbark, gum-topped box, spotted gum, grey gum, brush box, swamp mahogany, Queensland blue gum.
<b>Expected native pasture composition</b>	
Preferred	Black speargrass, Queensland bluegrass, kangaroo grass, hooky grass, leafy panic.
Non-indicator	Slender chloris, slender rat’s tail grass.
Non-preferred	Wiregrasses (e.g. dark), purple lovegrass, reed grass.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, green panic, Gatton panic, digit grass, tall finger grass, shrubby stylo, leucaena on deeper soils.

\* Denotes non-native ‘Expected Pasture Composition’ species.

## Blue gums on cracking clay



<b>Land resource area</b>	Floodplains.
<b>Landform</b>	Broad, low sloping floodplains on valley floors.
<b>Woody vegetation</b>	Tall open forest of Queensland blue gum and Moreton Bay ash with occasional broad-leaved apple, silver-leaved ironbark, rough-barked apple and broad-leaved ironbark. Understorey usually absent.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, hairy panic.
Non-indicator	Spring grass, liverseed (urochloa) grass, bamboo speargrass, umbrella grass.
Non-preferred	Wiregrasses (e.g. dark), slender chloris.
Legumes	Woolly glycine, rhynchosia, glycine pea, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, Angleton grass, leucaena, butterfly pea, Caatinga stylo, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Blue gums on granite



<b>Land resource area</b>	Granite hills.
<b>Landform</b>	Undulating rises to rolling hills.
<b>Woody vegetation</b>	Open forest to woodland of Queensland blue gum, silver-leaved ironbark and narrow-leaved ironbark. Understorey of wattle and minor beefwood.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, red natal grass*, silky umbrella grass, native millet.
Non-indicator	Pitted bluegrass grass, bottlewasher grasses, slender chloris, barbwire grass.
Non-preferred	Golden beard grass, comet grass.
Legumes	Emu foot, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, fine stem stylo, shrubby stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Blue gum on loam and duplex



<b>Land resource area</b>	Floodplains, Terraces.
<b>Landform</b>	Levees and levee backslopes along major streams and rivers, and the upper slopes of gently undulating relict alluvial plains and high terraces.
<b>Woody vegetation</b>	Tall open forest to woodland of Queensland blue gum and Moreton Bay ash or Queensland blue gum and rough-barked apple with occasional silver-leaved ironbark and narrow-leaved ironbark. Understorey usually absent.
<b>Expected native pasture composition</b>	<b>Southern black speargrass pastures.</b>
Preferred	Black speargrass, Queensland bluegrass, kangaroo grass, hooky grass, leafy panic.
Non-indicator	Slender chloris, slender rat's tail grass.
Non-preferred	Wiregrasses (e.g. dark), purple lovegrass, reed grass.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, digit grass, tall finger grass, shrubby stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Box on clay



<b>Land resource area</b>	Floodplains.
<b>Landform</b>	Backplains and levee backslopes of alluvial plains and creek flats.
<b>Woody vegetation</b>	Tall open woodland to open forest of poplar box, gum-topped box, broad-leaved apple. Understorey often absent.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, paspalum*.
Non-indicator	Pitted bluegrass, barbwire grass.
Non-preferred	Wiregrasses (e.g. dark).
Legumes	Woolly glycine, rhynchosia, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, Angleton grass, Bambatsi panic, leucaena, butterfly pea, Caatinga stylo, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Box on erosive soils



<b>Land resource area</b>	Terraces.
<b>Landform</b>	Broad low sloping, higher lying relict alluvial plains.
<b>Woody vegetation</b>	Open forest or woodland of poplar box, narrow-leaved ironbark, gum-topped box and occasionally rusty gum. Understorey usually absent, occasionally wilga and wattles.
<b>Expected native pasture composition</b>	Wiregrass – pitted bluegrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop.
Non-indicator	Spider grass, bottlewasher grasses, silky umbrella grass.
Non-preferred	Wiregrasses (e.g. dark), slender chloris.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, shrubby stylo.

\* Denotes non-native 'Expected Pasture Composition' species.

## Brigalow and brigalow belah



<b>Land resource area</b>	Undulating plains, Relict alluvial plains.
<b>Landform</b>	Gently undulating relict alluvial plains and higher lying level plains, and most slope positions on undulating low rises (slopes 1–4%).
<b>Woody vegetation</b>	Brigalow and brigalow belah open forest in association with wattles, wilga and softwood scrub.
<b>Expected native pasture composition</b>	Brigalow pastures.
Preferred	Brigalow grass, Queensland bluegrass, hooky grass, leafy panic.
Non-indicator	Slender chloris, slender rat's tail grass.
Non-preferred	Dark wiregrass, purple lovegrass.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Green panic, Gatton panic, creeping bluegrass, Angleton grass, Rhodes grass, buffel grass, Caatinga stylo, butterfly pea, siratro, leucaena, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Brigalow with melonholes



<b>Land resource area</b>	Relict alluvial plains.
<b>Landform</b>	Higher lying level plains, and mid slopes and crests of broad low rises.
<b>Woody vegetation</b>	Brigalow, black tea tree belah open forest.
<b>Expected native pasture composition</b>	Brigalow pastures.
Preferred	Brigalow grass, Queensland bluegrass, silky browntop.
Non-indicator	Native millet, spring grass, umbrella canegrass, slender chloris.
Non-preferred	Wiregrasses (e.g. dark), tall chloris.
Legumes	Woolly glycine, rhynchosia.
<b>Suitable sown pastures</b>	Creeping blue grass, Rhodes grass, green panic, buffel grass, Angleton grass, Bambatsi panic, Caatinga stylo, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Gum-topped box



<b>Land resource area</b>	Terraces and Relict alluvial plains.
<b>Landform</b>	Slightly elevated level to gently undulating relict floodplains, backplains and slightly higher terraces of major streams.
<b>Woody vegetation</b>	Open forest to woodland of gum-topped box, narrow-leaved ironbark and poplar box, with scattered rusty gum and Queensland blue gum. Occasional understorey of wattles, myrtle tree and beefwood.
<b>Expected native pasture composition</b>	Wiregrass – pitted bluegrass pastures.
Preferred	Black speargrass, forest bluegrass, barbwire grass, kangaroo grass, pitted bluegrass grass.
Non-indicator	Spider grass (native couch), bottlewasher grasses, umbrella grass.
Non-preferred	Wiregrasses (e.g. dark), slender chloris.
Legumes	Woolly glycine, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	None suitable.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark & bloodwood on non-cracking clay



<b>Land resource area</b>	Volcanic uplands.
<b>Landform</b>	Undulating rises and mid to lower slopes of low hills and ranges.
<b>Woody vegetation</b>	Woodlands of silver-leaved and narrow-leaved ironbarks and variable-barked bloodwood with occasional Queensland blue gum and areas of softwood scrub. Understorey usually absent.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, paspalum*.
Non-indicator	Pitted bluegrass, Indian couch*, barbwire grass, silkyheads.
Non-preferred	Wiregrasses (dark, erect kerosene), slender chloris, woodland lovegrass.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, Gatton panic, Caatinga stylo, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark & spotted gum on duplex and loam



<b>Land resource area</b>	Uplands sediments.
<b>Landform</b>	Gently to moderately inclined, undulating plains to slopes and rises of low hills (slopes 3–12%), with areas of steep hills (up to 40%).
<b>Woody vegetation</b>	Woodland to open forest of narrow-leaved ironbark and spotted gum, occasionally bloodwood, with an understorey of wattle and whitewood.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, paspalum*.
Non-indicator	Pitted bluegrass, Indian couch*, barbwire grass, silkyheads.
Non-preferred	Wiregrasses (e.g. dark, erect kerosene), woodland lovegrass, slender chloris.
Legumes	Woolly glycine, emu foot, creeping tick trefoil, rhynchosia.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, shrubby stylo.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark on basalt upper slopes and benches



<b>Land resource area</b>	Basalt rises.
<b>Landform</b>	Upper slopes and crests, including those on ridges, of undulating to low rises to steep hilly terrain and stony knolls.
<b>Woody vegetation</b>	Woodland or open forest of silver-leaved and/or narrow-leaved ironbarks in association with Queensland blue gum, variable-barked bloodwood, mountain coolibah and Moreton Bay ash. Understorey is usually absent.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, hairy panic.
Non-indicator	Spring grass, liverseed (urochloa) grass, bamboo speargrass, umbrella grass.
Non-preferred	Wiregrasses (e.g. dark), slender chloris.
Legumes	Rhynchosia, creeping tick trefoil, glycine pea, woolly glycine.
<b>Suitable sown pastures</b>	Rhodes grass, panic (green), creeping bluegrass, buffel grass, Angleton grass, Caatinga stylo, Desmanthus.

\* Denotes non-native 'Expected Pasture Composition' species.

## Mixed open forest on duplex and loam



<b>Land resource area</b>	Basalt rises, Volcanic uplands, Terraces, Relict alluvial plains, Red tablelands.
<b>Landform</b>	Widespread occurrence on mid, lower and upper slopes and crests of low basalt rises and stony knolls; upper slope positions on relict alluvial plains; mid and lower slopes of undulating plains and low hills, and mid to upper slopes of broad rises.
<b>Woody vegetation</b>	Open forest or woodland of gum-topped box, silver-leaved ironbark, narrow-leaved ironbark, with occasional Queensland blue gum, broad-leaved apple, pink bloodwood and spotted gum. Scattered occurrences of rusty gum, and wattle and dogwood.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, Queensland bluegrass, kangaroo grass, hooky grass, leafy panic.
Non-indicator	Slender chloris, slender rat's tail grass.
Non-preferred	Wiregrasses (e.g. dark), purple lovegrass, reed grass.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, shrubby stylo.

\* Denotes non-native 'Expected Pasture Composition' species.

## Narrow-leaved ironbark on granite



<b>Land resource area</b>	Granite hills.
<b>Landform</b>	Undulating rises to rolling hills.
<b>Woody vegetation</b>	Open forest to woodland of narrow-leaved ironbark, silver-leaved ironbark and Queensland blue gum and wattles.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, red natal grass*, barbwire grass, silky umbrella grass, hairy panic.
Non-indicator	Pitted bluegrass grass, niggerheads, bottlewasher grasses, woodland lovegrass.
Non-preferred	Wiregrasses (e.g. dark, erect kerosene), reed grass, golden beard grass.
Legumes	Woolly glycine, emu foot, glycine pea.
<b>Suitable sown pastures</b>	Oversow with legumes: fine stem stylo, shrubby stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Narrow-leaved ironbark and wattles



<b>Land resource area</b>	Ranges.
<b>Landform</b>	Crests and slopes of steep hills and mountains.
<b>Woody vegetation</b>	Woodland to open forest of narrow-leaved ironbark, silver-leaved, bloodwood, and spotted gum. If understorey present often wattles, rosewood, whitewood or beefwood.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, barbwire grass, pitted bluegrass, native oatgrass, kangaroo grass.
Non-indicator	Many-headed grass, kerosene grass, bottlewasher grasses.
Non-preferred	White speargrass.
Legumes	Narrow-leaved indigo, glycine pea.
<b>Suitable sown pastures</b>	Oversow with shrubby stylo.

\* Denotes non-native 'Expected Pasture Composition' species.

## Silver-leaved ironbark on cracking clay



<b>Land resource area</b>	Basalt rises.
<b>Landform</b>	Undulating plains to rolling hills.
<b>Woody vegetation</b>	Open forest or woodland of silver-leaved ironbark and Queensland blue gum with occasional narrow-leaved ironbark, variable-barked bloodwood. Understorey is usually absent.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, forest bluegrass, Queensland bluegrass, scentedtop, paspalum*.
Non-indicator	Native millet, hairy panic, barbwire grass, slender chloris.
Non-preferred	Wiregrasses (e.g. dark).
Legumes	Woolly glycine, rhynchosia, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, Caatinga stylo, Desmanthus, leucaena on deep soils.

\* Denotes non-native 'Expected Pasture Composition' species.

## Silver-leaved ironbark on granite



<b>Land resource area</b>	Granite hills.
<b>Landform</b>	Undulating rises with broad hill crests on granite.
<b>Woody vegetation</b>	Open forest to woodland of silver-leaved ironbark, narrow-leaved ironbark and Queensland blue gum. Understorey of wattles and minor beefwood.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, red natal grass*, barbwire grass.
Non-indicator	Pitted bluegrass grass, many-headed wiregrass, silky umbrella grass, feathertop Rhodes grass*.
Non-preferred	Dark wiregrass, reed grass, golden beard grass.
Legumes	Rattlepods, Birdsville indigo, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, fine stem stylo, shrubby stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Softwood scrub



<b>Land resource area</b>	Undulating plains, Red tablelands.
<b>Landform</b>	Mid to upper slopes and crests of gently sloping remnant plateaus and near scarp margins; slopes below scarps; and low hills adjacent to plateau remnants.
<b>Woody vegetation</b>	Softwood scrub (vines, bottle trees, white cedar, crows ash, figs) and open forest to open woodland of narrow-leaved ironbark in association with Queensland blue gum, blackbutt, spotted gum, Gympie messmate, grey gum, tallowwood or Yarraman ironbark depending on landscape position. Occasional bloodwoods, rusty gums, she-oaks and silver-leaved ironbarks with an understorey of wattles, red ash and dogwood.
<b>Expected native pasture composition</b>	
Preferred	Green panic*.
Non-indicator	
Non-preferred	Wiregrasses.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Green panic, Rhodes grass, buffel grass, Gatton panic, digit grass, tall finger grass, shrubby stylo, Caatinga stylo, Wynn cassia, siratro, leucaena.

\* Denotes non-native 'Expected Pasture Composition' species.

## Spotted gum ridges



<b>Land resource area</b>	Ranges.
<b>Landform</b>	Crests and hillslopes of undulating rises to low hills to mountains.
<b>Woody vegetation</b>	Spotted gum open forest or woodland frequently associated with narrow-leaved ironbark. Other species that may occur include bloodwoods, rusty gum, and gum-topped box. An understorey may include red ash, currant bush, grevilleas and wattles.
<b>Expected native pasture composition</b>	Wiregrass – pitted bluegrass pastures.
Preferred	Black speargrass, barbwire grass, pitted bluegrass, native oatgrass, kangaroo grass.
Non-indicator	Erect kerosene grass, kerosene grass, silkyheads.
Non-preferred	Dark wiregrass, five minute grass, comet grass.
Legumes	Rattlepods, glycine pea.
<b>Suitable sown pastures</b>	None suitable.

\* Denotes non-native 'Expected Pasture Composition' species.

## Tall open forest of snuffy soils



<b>Land resource area</b>	Red tablelands.
<b>Landform</b>	Upper slopes and crests of plateau remnants and some low rises.
<b>Woody vegetation</b>	Closed softwood scrub associated with open forest of narrow-leaved ironbark, grey gum, tallowwood, Gympie messmate and Yarraman ironbark with occasional bloodwoods, spotted gums and understorey of wattles and red ash.
<b>Expected native pasture composition</b>	Southern black speargrass pastures.
Preferred	Black speargrass, Queensland bluegrass, kangaroo grass, hooky grass, leafy panic.
Non-indicator	Slender chloris, slender rat's tail grass.
Non-preferred	Wiregrasses (e.g. dark), purple lovegrass, reed grass.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, green panic, digit grass, tall finger grass, leucaena, shrubby stylo, Caatinga stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Bloodwood and stringybark (coastal plains)



<b>Land resource area</b>	Coastal plains (Glanville <i>et al</i> 1991).
<b>Landform</b>	Level to gently undulating plains and low hills.
<b>Woody vegetation</b>	Bloodwoods, stringybarks, narrow-leaved ironbark, grey ironbark, Queensland peppermint and smooth-barked apple.
<b>Expected native pasture composition</b>	
Preferred	Barbwire grass, black speargrass, kangaroo grass.
Intermediate	Queensland blue couch*, cockatoo grass.
Non-preferred	Poverty grass, blady grass
Legumes	Rhynchosia, woolly glycine, glycine pea, Desmodium sps.
<b>Suitable sown pastures</b>	Rhodes grass, creeping bluegrass, signal grass, pangola grass, lotononis, shrubby and Caribbean stylos, siratro, villomix.

\* Denotes non-native 'Expected Pasture Composition' species.

## Blue gum flats



<b>Land resource area</b>	Alluvium (Glanville <i>et al</i> 1991).
<b>Landform</b>	Level alluvial plains (moderately extensive).
<b>Woody vegetation</b>	Blue gum, Moreton bay ash, silver-leaved ironbark, gum-topped box.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, scentedtop, Queensland bluegrass, black speargrass, paspalum*, kangaroo grass.
Intermediate	Pitted bluegrass, Queensland blue couch*, Angleton grass*, bahia grass*.
Non-preferred	Wiregrasses, blady grass, swamp foxtail.
Legumes	Rhynchosia, woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, pangola grass, fine stem, shrubby and Caribbean stylos, siratro, lotononis.

\* Denotes non-native 'Expected Pasture Composition' species.

## Blue gum, ironbark & bloodwood slopes & hollows



<b>Land resource area</b>	Granite (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating to rolling rises and plains.
<b>Woody vegetation</b>	Blue gum, narrow-leaved ironbark, bloodwood and wattles.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, tambookie grass, black speargrass, kangaroo grass.
Intermediate	Pitted bluegrass, Queensland blue couch*, barbwire grass, thatch grass*, bahia grass*.
Non-preferred	Wiregrasses, bottlewasher grasses, rat's tail grasses.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes, signal, pangola grass, fine stem and shrubby stylos, siratro, lotononis, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Gum-topped box



<b>Land resource area</b>	Uplifted coastal plains, metamorphic (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating plains, low hills and ridges.
<b>Woody vegetation</b>	Gum-topped box, narrow-leaved ironbark, blue gum, spotted gum and wattles.
<b>Expected native pasture composition</b>	
Preferred	Pitted bluegrass, barbwire grass, kangaroo grass.
Intermediate	Queensland blue couch*, lovegrasses.
Non-preferred	Wiregrasses.
Legumes	Woolly glycine, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Not suitable for sown pastures. Oversow with legumes: shrubby stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Hoop pine scrub



<b>Land resource area</b>	Metamorphic (Glanville <i>et al</i> 1991).
<b>Landform</b>	Mountains and undulating to rolling rises and plains.
<b>Woody vegetation</b>	Hoop pine with depauperate rainforest understorey. Crow's ash, Burdekin plum and bottletrees also occur.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, Queensland bluegrass, black speargrass, kangaroo grass.
Intermediate	Pitted bluegrass, tambookie grass, Queensland blue couch*, Angleton grass*.
Non-preferred	Wiregrasses, native chloris.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, fine stem and shrubby stylos, siratro, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark, stringybark and supplejack ridges



<b>Land resource area</b>	Granite (Glanville <i>et al</i> 1991).
<b>Landform</b>	Mountains and low hills.
<b>Woody vegetation</b>	Narrow-leaved ironbark, grey ironbark, white mahogany, white stringybark, thin-leaved stringybark, spotted gum, bloodwoods, turpentine, wattles, grass tree and supplejack.
<b>Expected native pasture composition</b>	
Preferred	Barbwire grass, black speargrass, kangaroo grass.
Intermediate	Pitted bluegrass, Queensland blue couch*, poverty grass.
Non-preferred	Wiregrasses, blady grass, rat's tail grasses.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Not suitable for sown pastures. Oversow with legumes: shrubby stylo, fine stem stylo, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark and bloodwood on non-cracking clay



<b>Land resource area</b>	Granite, metamorphic, acid volcanic (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating to rolling rises and plains.
<b>Woody vegetation</b>	Silver-leaved ironbark, narrow-leaved ironbark, bloodwood and wattles.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, Queensland bluegrass, black speargrass, kangaroo grass.
Intermediate	Pitted bluegrass, Queensland blue couch*, barbwire grass, tambookie grass.
Non-preferred	Wiregrasses, bottlewasher grasses, rat's tail grasses.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, signal grass, fine stem and shrubby stylos, siratro, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark and blue gum on basalt ridges



<b>Land resource area</b>	Basalt (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating plains, low hills and plateau remnants.
<b>Woody vegetation</b>	Blue gum, silver-leaved ironbark, Moreton bay ash, pink bloodwood.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, Queensland bluegrass, Angleton grass* (naturalised).
Intermediate	Pitted bluegrass, Queensland blue couch*.
Non-preferred	Wiregrasses.
Legumes	Rhynchosia, creeping tick trefoil, glycine pea, woolly glycine.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, Angleton bluegrass, Caatinga and fine stem stylos, siratro, leucaena.

\* Denotes non-native 'Expected Pasture Composition' species.

## Ironbark and spotted gum on duplex and loam



<b>Land resource area</b>	Granite, acid volcanic, metamorphic (Glanville <i>et al</i> 1991).
<b>Landform</b>	Mountains, low hills and minor undulating plains.
<b>Woody vegetation</b>	Spotted gum, narrow-leaved ironbark, grey ironbark, bloodwoods and wattles.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, tambookie grass, black speargrass, kangaroo grass.
Intermediate	Pitted bluegrass, Queensland blue couch*, barbwire grass, cockatoo grass, golden beard grass.
Non-preferred	Wiregrasses, blady grass, lovegrasses.
Legumes	Woolly glycine, rhynchosia, emu foot, creeping tick trefoil.
<b>Suitable sown pastures</b>	Creeping bluegrass, Rhodes grass, signal grass, fine stem and shrubby stylos, siratro, Wynn cassia.

\* Denotes non-native 'Expected Pasture Composition' species.

## Mixed eucalypts on uplifted coastal plains



<b>Land resource area</b>	Uplifted coastal plains (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating plains, low hills and ridges.
<b>Woody vegetation</b>	Bloodwoods, stringybarks, narrow-leaved ironbark, grey ironbark, red ironbark, Queensland peppermint, spotted gum, smooth-barked apple and grass trees.
<b>Expected native pasture composition</b>	
Preferred	Golden beard grass, barbwire grass, black speargrass, kangaroo grass.
Intermediate	Queensland blue couch*, cockatoo grass.
Non-preferred	Poverty grass, blady grass.
Legumes	Woolly glycine, rhynchosia, Desmodium sps.
<b>Suitable sown pastures</b>	Not suitable for sown pastures. Oversow with legumes: lotononis, shrubby and Caribbean stylos, siratro.

\* Denotes non-native 'Expected Pasture Composition' species.

## Softwood scrub



<b>Land resource area</b>	Basalt (major), Alluvium (minor) (Glanville <i>et al</i> 1991).
<b>Landform</b>	Undulating plains, low hills and plateau remnants.
<b>Woody vegetation</b>	Softwood scrub including bottle trees, white cedar and crow's ash with depauperate rainforest understorey.
<b>Expected native pasture composition</b>	
Preferred	Forest bluegrass, Queensland bluegrass and scentedtop (run down pasture). Green panic, Rhodes grass in new pastures.
Intermediate	Pitted bluegrass, Queensland blue couch*, Angleton grass*.
Non-preferred	Wiregrasses.
Legumes	Woolly glycine, glycine pea.
<b>Suitable sown pastures</b>	Green panic, Rhodes grass, pangola grass, leucaena, siratro, axilaris, glycine.

\* Denotes non-native 'Expected Pasture Composition' species.

## Tea tree flats



<b>Land resource area</b>	Alluvium (major); sandplain and coastal plain (minor) (Glanville <i>et al</i> 1991).
<b>Landform</b>	Level alluvial plains (moderately extensive).
<b>Woody vegetation</b>	Paperbark tea tree, bloodwoods, blue gum, swamp mahogany.
<b>Expected native pasture composition</b>	
Preferred	Golden beard grass, black speargrass, kangaroo grass.
Intermediate	Queensland blue couch*.
Non-preferred	Poverty grass.
Legumes	Woolly glycine, glycine pea, Desmodium sps.
<b>Suitable sown pastures</b>	Pangola grass, humidicola, lotononis, villomix.

\* Denotes non-native 'Expected Pasture Composition' species.



## Legumes for Burnett pastures

Select legumes based on their suitability for the soil type and their suitability for the production system. Table 4 lists the legumes recommended for the Burnett district. Descriptions are provided outlining the characteristics of each species.

### Recommended planting rates and inoculant

Recommended planting rates vary considerably. If a legume is used in a

mix of other legumes the lower rate would be used whereas the higher rate would be planted if the legume is being used as a single species planting. Adjust the planting rate if using coated seed (see Table 5).

Many tropical legumes are hardseeded. This characteristic enables the seed to remain in the soil for long periods and to germinate when conditions are favourable. Most hardseeded legumes are resistant to fire. To encourage germination of these seeds in a sown pasture it is often beneficial to 'scarify' or 'scar' the seed coat. Seed companies generally do this if necessary.

Table 4 – Legumes for Burnett soils


Soil	Suitable legumes
Red scrub	Leucaena, desmanthus, white clover (Haifa), lucerne, shrubby stylo (seca)
Black scrub	Wynn cassia, leucaena, desmanthus white clover (Haifa), lucerne, Caatinga stylo
Red forest	Wynn cassia, lucerne, leucaena, white clover, desmanthus, shrubby stylo
Brown forest	Lucerne, white clover, desmanthus, Wynn cassia, seca stylo, Caatinga stylo
Black forest (ironbark)	Lucerne, medics, desmanthus, white clover, leucaena, vetch, siratro, seca stylo, Caatinga stylo
Black alluvial (blue gum)	Lucerne, medics, white clover, desmanthus, Maku lotus, siratro, leucaena
Hardsetting (box trees )	Wynn cassia, seca stylo
Sandy surface (ironbark)	Seca stylo, Wynn cassia, siratro
Stony ridges (spotted gum)	Seca stylo, Wynn cassia
Coastal loams and clays	Wynn cassia, siratro, leucaena, Maku lotus, creeping vigna, joint vetch

Table 5 – Recommended planting rates for legumes

Species	Planting rate	Rhizobium inoculant
Annual medics	0.3–2 kg/ha in pastures. Up to 8 kg/ha for snail medic forage cropping.	Group AM
Burgundy bean	3–5 kg/ha. Seed coat needs scarifying.	CB1717
Butterfly pea	3–4 kg/ha	Tropical group M
Clover	0.3–1 kg/ha	Group B, TA1
Creeping vigna	0.5–3 kg/ha	Group I
Desmanthus	2 kg/ha. Seed coat needs scarifying.	CB3126
Forage peanuts	Main variety is Pinto which can be planted by seed or runners. Prine needs to be planted from runners.	CIAT 3101
Joint vetch	0.5–3 kg/ha	CB2312
Lab lab	12–20 kg/ha for single sowing with grass; 5–8 kg/ha if sown with other legumes.	Group J
Leucaena	2–2.5 kg/ha	Leucaena, CB81
Lotononis	0.1–0.25 kg/ha. Also established via runners or cattle dung.	Group L, CB376
Lucerne	1–6 kg/ha	Group A, CC169
Maku lotus	2–3 kg/ha	Group D, CC829
Siratro	0.5–2 kg/ha. Seed coat needs scarifying.	Group M
Serradella	8 kg/ha of podded seed.	Group G, WU425
Stylo, fine stem	2–5 kg/ha. Seed coat needs scarifying.	CB82, CB1650, CB1552
Stylo, shrubby	1–2 kg/ha. Seed coat needs scarifying.	Group S but will readily combine with native rhizobia
Vetch	2–4 kg/ha	Group E, SU391
Wynn cassia	0.3–1 kg/ha	Group M



photo: David Gramshaw



## Annual medics

### *Medicago spp*

Many types of medics have been trialled in the Burnett including barrel medic (*M. trunculata*), common burr medic (*M. polymorpha*) and snail medic (*M. scutella*). These are winter/spring annuals and grow well with early spring rainfall.

They are suited to clay or loam soils with a neutral to high pH. The snail medics such as Sava and Kelson can be used in crop-pasture rotations while the barrel medics such as Jemalong, Cyperus and Paraggio perform best in pastures. Autumn, after the threat of hot weather has passed, is the best time for planting.

Generally, snail medics have not persisted well in the Burnett, but burr medics are common.

Medics will do best in wet winters. Bloat can be a problem if there is a large bulk of these legumes. Barrel medics are a worthwhile addition to wheat crops to improve productivity and grazing on harvested stubble. Seed is generally not expensive and they have been found to be very beneficial for stock.

## Burgundy bean

### *Macroptilium bracteatum*

Burgundy bean is a summer-growing, twining perennial with dark red to burgundy flowers.

It was selected for use on clay soils. It is drought tolerant and has survived for up to three years on shallow clay soils.

It has the ability to germinate and grow earlier in the season than other tropical legumes. It is susceptible to frost.

It is very palatable to stock and needs spelling to allow it to go to seed and persist in pastures. Burgundy bean is well suited to sandy loams and clay loams. Tolerates slightly acid to alkaline soils.

Seeds are large and can germinate in less-than-ideal conditions. It can be hard seeded and the seed coat may require scarifying to ensure adequate germination. There are two varieties of burgundy bean that are marketed together: Cardarga is an upright growing plant; Juaninta has 19% protein.



## Butterfly pea

*Clitoria ternata*

Butterfly pea is a tropical summer-growing, twining perennial with mauve to blue flowers. It can vary in appearance depending on the growing conditions. Commonly used as an ornamental garden creeper.

Butterfly pea grows well on a variety of soil types but performs best on heavy alkaline black clay soils.

This tropical plant will not persist in cold areas subject to heavy frosts. In the Burnett it needs to be planted high in the landscape and be well established to survive winter frosts.

It is a very palatable, excellent quality fodder that is also suitable for hay.

Milgarra is the main variety. This variety has some tolerance to flooding and waterlogging.




## Clover

*Trifolium spp*

There are a number of clovers available but generally white clover has been the most successful in the Burnett.

Clovers are temperate legumes suited to high moisture situations, but are worth planting because they are productive in wet winters and early in spring. Clovers are the most persistent legume in kikuyu pastures.

The main variety is Haifa. There are many other varieties but availability varies. Choose a variety suited to the sub-tropics.



## Desmanthus

*Desmanthus virgatus*

Desmanthus is a summer-growing, woody perennial shrub. It usually grows to around 70 cm in height but can grow as tall as 1.5 m. Its taproot grows to a depth of 0.5 m. Desmanthus grows on a wide variety of soils but is best suited to alkaline duplex and cracking clay soils. Frost will defoliate the shrubs but it will re-grow from established crowns after frosting provided there is sufficient soil moisture.

The seed coat is very hard and established stands should be allowed to set seed for two years to build up seed reserves. Once established it will tolerate heavy grazing. Desmanthus is palatable to cattle but not as favoured as leucaena.

The three cultivars, Marc, Bayamo and Uman varieties, have short, mid and late season flowering respectively. Progardes is a selection of these three types.

Desmanthus has an average crude protein of 21%.



## Forage peanuts

*Arachis spp*

Forage peanuts have not been used widely in Burnett pastures because they are sensitive to cooler conditions. They may do well in protected situations.

There are two main varieties. Pinto grown from seed and is used in coastal areas as a horticultural ground cover. Prine is grown from cuttings and has been trialled in the dairy industry in Central Queensland.





### Joint vetch

*Aeschynomene americana* and  
*A. falcata*

American joint vetch is suited to wet soils of low fertility. It can grow to 2 m tall and is susceptible to frosts. Bargoo is a smaller variety that can handle drier and cooler conditions. Both plants are very palatable.



### Lab lab

*Lablab purpureus*

Lab lab is a summer growing annual that occasionally persists as a short-lived perennial. It is a vigorous robust trailing and twining plant. Stems grow to 3–6 m in length.

It is an ideal plant for grazing and cropping rotations or as an addition to forage crops. It performs well as a green manure crop and has very good forage quality for grazing or as hay.

Lab lab does not tolerate frost but will grow in a wide range of soils from deep sands to heavy clays, provided drainage is good and the pH is 4.5–7.5. Lab lab has an average crude protein of 26%.

There are two varieties. Rongai has white flowers and is a late-flowering, with high dry matter production. Highworth is an earlier-flowering variety originally intended for grain production. It has purple flowers and black seeds.

Lab lab is an excellent addition to summer fodder crops.



## Leucaena

### *Leucaena leucocephala*

Leucaena is a tree legume needing deep, fertile, well drained soil. Mature trees grows to 18 m high but should be kept lower in a grazing situation. It does best in areas with minimal frost. Leucaena is slow to establish and needs careful weed management in the early stages.

Leucaena is costly to establish but very productive. Well managed stands in the Burnett have remained productive for over 30 years. Leucaena can be toxic to cattle. Inoculate cattle grazing leucaena with a rumen bacteria to ensure digestibility of the plant.

There are three main varieties. Peru is one of the original varieties introduced by CSIRO. Cunningham is more highly branched and 30% higher yielding than Peru. Tarramba establishes more rapidly, and is marginally more psyllid-resistant and cool-tolerant than Peru and Cunningham. Redlands is a psyllid resistant variety.

Leucaena has potential to be an invasive weed if not managed correctly. The Leucaena Network has a Code of Practice for members to follow to avoid weed problems.

For more information on leucaena it is recommended to read the Meat & Livestock Australia publication *Leucaena: A guide to establishment and management*. The Leucaena Network also provides valuable information about establishing and managing plantations ([www.leucaena.net](http://www.leucaena.net)).



## Lotononis

### *Lotononis bainesii*

Lotononis is an early summer-growing perennial with moderate frost tolerance. It prefers lighter well drained soils but is found on a variety of soil types; gravelly, sandy and loamy soils, but not generally on clays.

It has yellow flowers and is a low growing, very palatable species, common on roadsides in the Burnett. It tolerates low fertility soils and heavy grazing. It is slow, and often difficult, to establish due to hard seeds, but persists once established.

Lotononis spreads by both runners and seed. It does not do well if it is shaded out by taller grasses particularly those with runners. It performs well under rotational grazing regimes. Lotononis has an average crude protein range of 9–25%.



## Lucerne

### *Medicago sativa*

Lucerne is the king of fodder crops. It is a perennial suited to fertile, deep, well drained soils. It does not persist well on shallow or acidic soils.

All lucerne cultivars are summer-growing, but some cultivars are more winter-active than others. As a result, cultivars can be described as winter-dormant, semi winter-dormant, winter-active, and highly winter-active.

Growth in semi-dormant and dormant cultivars slows down during the colder months. Generally these types have lower crowns and are more persistent in dryland grazing systems.

Highly winter-active cultivars will provide year-round feed with only a slightly reduced production in winter, especially if they are irrigated. Winter-active cultivars fall between these extremes. They have moderate winter growth and are moderately persistent under grazing.

Bloat can be a problem in pastures containing a high proportion of lucerne. Rotational grazing is essential to maintain plant population.

Choose a lucerne variety that suits your situation, there are many cultivars available bearing in mind the semi-dormant and dormant cultivars are best and most persistent for dryland pastures.



## Maku lotus

### *Lotus pedunculatus*

This is a perennial that is suited to waterlogged areas. It has a variable growth habit depending on the environment, growing up to 60 cm high.

This legume needs very good moisture levels but is tolerant of frost, heavy grazing and low fertility acid soils. It has poor drought tolerance but does not cause bloat.



photo: David Gramshaw

## Serradella

### *Ornithopus compressus*

Serradella is a winter/spring annual for deep, light well-drained soils and acidic sandy soils. It is a non-bloating legume. Serradella has not been planted widely in the Burnett but does have some potential in southern areas.

The Maderia and Santorini varieties of yellow serradella are recommended for southern Queensland.

Slender serradella, the Jebala variety, flowers mid-season and is hardseeded. It can tolerate poorly drained soils (shallow, stony soils that can become very wet) better than yellow serradella, and can produce a good bulk of forage in late spring.



## Siratro

*Macroptilium atropurpureum*

Siratro is a perennial vine with a deep, swollen taproot and trailing, climbing and twining stems.

It grows on a wide range of soils from dark cracking clays to sands and gravels, but does best on more fertile soils. It is sensitive to frost and rust but is drought resistant.

Siratro persists best under rotational grazing and requires spelling to set seed to ensure persistence. Siratro seed requires scarification before planting. It has an average crude protein of 25%. The Aztec variety is rust resistant.



## Stylo, Caatinga

*Stylosanthes seabrana*

Caatinga stylo differs from the other commercially available stylos in that it grows well on clay and loam soils. It is perennial and can be slow to establish in the first season but is particularly persistent and hardy with sown grasses including buffel grass. It has moderate frost and drought tolerance.



### Stylo, fine stem

*Stylosanthes guianensis* var.  
*intermedia*

Fine stem stylo is a low growing (up to 30 cm high) perennial with fine hairs visible on the stems. It is tolerant of cold, fire and heavy grazing. It needs deep, free draining sandy soils to perform best.

Fine stem stylo is palatable to stock and responds well to regular grazing. It may be lost from a pasture if it is shaded out by tall grasses. Its resistance to fire makes fine stem stylo a useful addition to native pastures. Cattle will spread the seed in their dung once it is established.

It is hard seeded so seed requires scarification before planting. Establishment is usually slow. Fine stem stylo has an average crude protein of 16%.



### Stylo, shrubby

*Stylosanthes scabra*

Shrubby stylo is a perennial shrub legume that grows to 2 m tall with a long taproot. It is tolerant of heavy grazing and drought. Plants are slow to grow in the first year and can become very woody with age.

Shrubby stylo is sensitive to heavy frosts and waterlogging. It has hard seeds so establishment is slow if the seed is not scarified. Seed will spread over time in manure of stock and in pods attached to hair.

The two varieties, Seca and Siran, are suited to a wide range of soils except for heavy clays. Caatinga stylo is more suited to heavier clay soils.



## Vetch

*Vicia villosa ssp dasycarpa*

Vetch is a spring growing annual suited to most soils except those susceptible to waterlogging and those that set very hard. It is a good pioneer species but needs to be spelled to allow seed to set. There is a local native spurred vetch that is common in Burnett pastures.

The main varieties are Namoi, Woolly pod and Poppany.



## Wynn cassia

*Chamaecrista rotundifolia*

Wynn cassia is a tropical legume that is quick to establish and seeds well. It often dies in winter and returns from seed. It is susceptible to frost and drops leaves under drought conditions.

It prefers lighter soils and can withstand heavy grazing. Palatability is a problem in some areas and pastures must be managed carefully to ensure grass is not overgrazed. Wynn cassia has the potential to form a pure sward leaving no grasses if the pasture is overgrazed. It has been used to make hay in some coastal areas.



Select grasses based on their suitability for the soil type and their suitability for the production system. Table 6 lists the grasses recommended for the Burnett district. Descriptions are provided outlining the characteristics of each species.

The best mix of pasture will depend on cost and availability of seed. The planting rates given vary, with the higher rates for single species planting and lower rates for mixes of species. Rates are for uncoated seed.

*Table 6 – Introduced grasses for the Burnett*

Soil	Tree type	Pasture grass
Red scrub	Scrub	Rhodes grass (Callide & Katamboora types), creeping blue grass, kikuyu, digit grass, panic, paspalum, brachiaria, setaria (Narok & Splenda)
Black scrub	Scrub	Rhodes grass (Callide & Katamboora types), creeping blue grass, kikuyu, digit grass, panic, paspalum, brachiaria, bambatsi, panic
Red forest	Narrow leaf ironbark	Rhodes grass, creeping blue grass, digit grass, kikuyu, paspalum, pangola
Brown forest	Narrow leaf ironbark	Rhodes grass, creeping blue grass, digit grass, paspalum, pangola
Black forest	Broad leaf ironbark	Rhodes grass, creeping blue grass, setaria (Narok & Splenda), bambatsi, paspalum, Swann blue grass, pangola, digit grass
Black alluvial	Blue gum	Rhodes grass, kikuyu, bambatsi, paspalum, Floren blue grass, setaria (Narok & Splenda)
Hardsetting duplex	Gum top box	Rhodes grass, creeping blue grass, digit grass, Indian blue grass (Medway)
Sandy surface	Ironbark	Creeping blue grass, Rhodes grass, digit grass, pangola, Indian blue grass (Medway)
Stony ridge	Spotted gum, wattles	Rhodes grass, creeping blue, digit grass, Indian blue grass (Medway)

## Recommended planting rates

Table 7 – Recommended planting rates for grasses

Grass	Planting rate	Comments
Bambatsi	2–4 kg/ha	seeds have long dormancy
Buffel grass	1–4 kg/ha	fluffy seed
Blue grass (creeping)	1–4 kg/ha	fluffy seed
Blue grass (Floren)	2–4 kg/ha	suited to alkaline soils
Blue grass (Indian)	3–4 kg/ha	suited to low fertility soils
Blue grass (Swann forest)	2 kg/ha	fluffy seed
Brachiaria	8–10 kg/ha	Establishes well within 2 months
Digit grass	2–4 kg/ha	can spread by stolons
Kikuyu	0.5–1 kg/ha	spreads by runners
Panic	3–5 kg/ha	shade tolerant
Pangola	n/a	propagate via runners only
Paspalum	2–4 kg/ha	sticky seeds spread by animals
Purple pigeon grass	2–4 kg/ha	seeds have long dormancy
Rhodes grass	2–4 kg/ha	fluffy seed, most spread by runners
Sabi grass	1–4 kg/ha	seed readily available
Silk sorghum	2–4 kg/ha	responds to warmer soil temperature



## Bambatsi

*Panicum coloratum* var.  
*makarikariense*

Bambatsi has distinctive bluish leaves with a white mid-rib. The foliage grows to a height of 70 cm. Bambatsi spreads by seeds and stolons.

It is slow to establish, but tolerates light frosts and is very persistent once established. It is well adapted to heavy, self-mulching black clay soils, black soil creek flats and melon hole country, it is tolerant of temporary waterlogging, flooding, drought and moderate soil salinity.

Bambatsi is cold tolerant and has high forage quality. It has a lower nitrogen requirement than green panic. Seeds can have a long dormancy period. Photosensitization has occurred in sheep, goats, cattle and horses grazing Bambatsi but it is rare. Bambatsi is very palatable with crude protein ranging from 5–19%.



## Buffel grass

*Cenchrus ciliaris*

Buffel grass is very widely planted in central and western Queensland. It requires fertile soils and responds quickly to rain with excellent production. Plants will grow to 2 m high depending upon cultivar.

It is drought tolerant but can be slow to establish. Fresh seed has a period of dormancy. The seed is fluffy and difficult to sow in conventional seeding equipment unless pelleted. Buffel grass is not generally adapted to southern areas of the Burnett but will grow more widely in the northern areas. It is often worthwhile planting buffel grass on elevated red soils in the South Burnett. Buffel grass has an average crude protein of 6–16%. If it is their sole diet, buffel grass can cause a disease called ‘bighead’ in horses.

There are many varieties but the main ones grown in the Burnett are Biloela (the main tall variety now available), Gayndah (more prostrate and dense tillered) and American/USA (similar to Gayndah but has a longer flowering season).



## Blue grass (creeping)

*Bothriochloa insculpta*

Creeping blue grass is a perennial, stoloniferous tussock grass that grows to 0.9 m tall. The seed and leaves have a strong scent when crushed. It grows well on low fertility soils, but not on waterlogged soils or very heavy clays.

When growing on black soil creek flats creeping bluegrass can be unpalatable. Its main growth occurs in summer/autumn. It is tolerant of heavy grazing on most soils and has been cut for hay in the Burnett. There is slow regrowth in spring.

Seed is fluffy and difficult to spread unless coated. It generally takes more than one season to establish. Average crude protein levels range from 10% at the beginning of the growing season to 5% at the end of the season. Creeping blue grass is useful for erosion control.



The varieties suited to the Burnett are Hatch and Bisset. Hatch is long stemmed and more robust than the other varieties, runners do not root down. Bisset sends roots down from runners, is finer stemmed and flowers later than the other varieties. It is more palatable than Hatch.



### Blue grass (Floren)

*Dicanthium aristatum*

Floren blue grass is a selected line of the introduced species angleton grass which is common in Queensland. Floren blue grass is not widely planted in the Burnett at present.

It is a late maturing, palatable perennial growing to a height of 1.8 m with a vigorous, spreading growth habit. Floren blue grass is suited to alkaline, cracking clay soils.

On the Darling Downs it is used extensively to compete with the weed lippia. Floren blue grass will tolerate flooding and saline conditions. It is palatable after frosting and persists well under heavy grazing.



### Blue grass (Swann forest)

*Bothriochloa bladhii* subsp. *glabra*

Swann forest blue grass is a perennial tussock grass growing to 80 cm, generally with unbranched stems. The seedhead is green to purplish that, along with the leaf, has a distinctive odour when crushed. Swann forest blue grass makes good hay if it is cut before flowering but loses quality after that point. The main flowering is late in the season.

This sub species of blue grass has its origins in India and is different to Burnett blue grass, or native forest blue grass, *Bothriochloa bladhii*.

Swann forest blue grass grows well on both fertile and infertile soils but not on soils with high aluminium levels such as the red acid soils. The seed is fluffy and therefore difficult to sow. Maximum germination is reached 6–7 months after maturation. It has an average crude protein level range of 7–14%. It is susceptible to leaf rust and not as palatable as some other summer grasses.

Swann forest blue grass is more widely adapted than the native forest blue grass. It is palatable, tolerates heavy grazing and survives seasonal frosting.



## Blue grass (Indian)

*Bothriochloa pertusa*

Indian blue grass is a creeping stoloniferous and/or tufted perennial, commonly with pink to red stolons and upright stems. It grows to 50 cm before flowering and is 90 cm at maturity.

The leaf blades are greyish green and aromatic when crushed. The seedhead is purplish.

This grass is not widely planted in the Burnett at present however it is very common in northern Queensland where it has replaced speargrass. It is less palatable than many native grasses. It is generally unpalatable after flowering but is eaten readily when young and green.

Indian blue grass tolerates a wide range of soil types but should only be considered for poorer soils where options for other sown grasses are limited. It is a very successful coloniser due to its tolerance of low fertility soil and heavy grazing. This grass can produce significant quantities of seed.

In some areas it is considered a weed of pastures, although its presence is usually a symptom of declining fertility and excessive grazing. When young this grass can have a crude protein level up to 12% but when averaged with flowering plants the average crude protein drops to around 3–4%.

There are three varieties of Indian blue grass. Medway was selected for leaf and length of flowering. It is a more robust type, best adapted to lower rainfall areas. Keppel is late flowering with leafy runners and fine stems. Dawson is low growing and used as a turf grass.



## Brachiaria hybrids

### *Brachiaria crosses*

This hybrids are well-adapted to acid and neutral soils of moderate to low fertility. They tolerate high soil aluminium levels. This grass is persistent under seasonally dry conditions. It is productive, and capable of sustaining high stocking rates and grazing pressures. It has high nutritional value for ruminants. Responds well to nitrogen fertiliser.



## Digit grass

### *Digitaria milanijana*

Digit grass grows to 1.5 m. It is suited to low fertility soils but may need phosphorus and potassium on less fertile soils. It is very palatable, has early growth in spring and is drought hardy. It can spread by stolons. It may not establish well on heavy soils. Average crude protein ranges from 8 to 12%. There are four varieties available. Jarra grows to 1.8 m with broad dark green to purple leaves. Jarra does better in higher rainfall areas. Strickland grows to 1.5 m and has grey green leaves. It is more drought tolerant than Jarra.

Premier Digitaria has performed well with legumes and nitrogen on infertile soils, but it loses productivity when nitrogen declines.



## Kikuyu

### *Pennisetum clandestinum*

Kikuyu is a productive and useful grass that spreads by runners. It can be established from seed but is expensive and slow. Kikuyu usually grows 30–40 cm high and will form a tight mat if heavily grazed. It requires highly fertile red and black soils with adequate moisture and will tolerate low pH (acid) soils. It will respond well to high levels of nitrogen and phosphorous.

Kikuyu performs best if grazed to maintain a height no lower than about 5 cm and no higher than 15 cm. Kikuyu can become stemmy resulting in reduced leaf production if allowed to grow too tall. Rest kikuyu pastures after grazing down to 5 cm. Kikuyu can dominate and exclude other pasture grasses and legumes.

In high rainfall areas a soil borne parasite can cause 'kikuyu yellows'. Rust and various insects can also cause damage and production losses to kikuyu stands.

Crude protein levels can reach 25% in well fertilised stands. New growth is very palatable for stock and stock will often eat out small patches of kikuyu in a mixed pasture. Lactating dairy cows eating a high percentage of kikuyu in their diet may need calcium and sodium supplements. Nitrate poisoning, bloat and oxalate poisoning can also occur. Despite this kikuyu can provide valuable green feed in winter.

The two varieties of kikuyu are Whittet and Noonan. Whittet is a taller variety with broad leaves. It persists well on low fertility soils. Noonan seed is available but this variety is more suited to turf. It is resistant to kikuyu yellows.



## Panics

*Megathyrsus maximus*  
(was *Panicum maximum*)

The panics are a large group of grasses that are very variable in appearance. Green panic and Gatton panic are the most common cultivars with Hamil grass being one of the tall types grown more in the tropics.

Panic is a bright green tussock grass with a wide leaf and stout stems growing to 1.5 m tall. Green and Gatton panic can flower throughout the season. They are shade tolerant and respond quickly to spring rain often being the first grass to appear after long dry periods.

Panics are able to grow in a wide range of soil types but generally only persist on more fertile scrub soils. Being very palatable to stock it is easily grazed out of pastures that are not rested.

They respond well to fertilising, and will be replaced by other grasses as nitrogen becomes locked up in older pastures. Panic spreads well particularly in areas where it is not under grazing pressure.

Green panic is a fine stemmed Guinea grass (*Megathyrsus maximum* var. *trichoglume*) with soft leaves. It grows to 1.5 m tall and is shade tolerant. It has hairy stems in comparison to Gatton panic which has smooth stem nodes. Gatton panic is more vigorous than green panic with longer and wider leaves. It often has red to purple colouration on the stems and is easier to establish and manage.



## Pangola grass

*Digitaria eriantha*

Pangola grass is extremely variable in form. It is a perennial that spreads from runners. Only a very limited number of seeds are viable. Runners need to be harvested and planted into prepared soil during damp weather to propagate this grass.

Pangola grass is very productive on lighter soils but will grow on a large range of soil types. This grass competes well with African lovegrass and other weeds. It has a moderate tolerance to salinity, it is palatable and tolerates heavy grazing. Pangola grass responds well to fertiliser and is cut for hay in tropical areas. The average crude protein of pangola grass is 9–20%.



## Paspalum

*Paspalum dilatatum*

Paspalum is a leafy, tufted, sod-forming perennial, arising from short rhizomes 4–8 mm in diameter. Paspalum has a thick fibrous root system growing to more than 1 m deep. It is found throughout the Burnett on a wide range of moderately fertile soils but it does best on more fertile soils.

Paspalum responds well to fertiliser and provides stable ground cover to reduce the risk of water erosion. It is palatable, persistent and tolerates waterlogging.

The sticky seeds spread by attachment to animals and vehicles. An ergot (fungus) on seedheads can be a problem but paspalum rarely forms a high proportion of pastures in the Burnett and so there is insufficient ergot to cause ill-effects in cattle. The average crude protein ranges from 4–23%.

The Hi-Gane variety is suitable for wetter areas.



## Purple pigeon grass

*Setaria incrassata*

Purple pigeon grass forms a tussock growing to 2 m high. The seedhead is a distinctive spike. It is adapted to high fertility black and red clay soils where it establishes more reliably than Rhodes or panic grasses. Purple pigeon grass mixes well with lucerne and medics.

Purple pigeon grass is affected by frosts but recovers to give good spring and summer growth. It is best planted as a pure stand as it can be unpalatable and will not be grazed when there is other pasture available. It is most palatable when it is short and is useful for hay. Purple pigeon grass establishes quickly and could be suited to short term pasture leys on black soils. It is not recommended for horses due to potential oxalate poisoning. The seed can lay dormant for up to a year.



## Rhodes grass

*Chloris gayana*

Rhodes grasses are very common and widely planted in the Burnett area. Most varieties spread by runners but are also easily established from seed although the seed is fluffy and if uncoated, is difficult to sow using conventional planting equipment. It grows to 2 m tall and has roots to 4 m depth. Rhodes grass does best on fertile soils but will survive on lighter soils. It will respond well to fertiliser.

Rhodes grass has good tolerance to saline conditions and provides good ground cover. It has a crude protein range from 17% when green to 3% when dry. Palatability reduces markedly after drying out or frosting.

There are several varieties of Rhodes grass. Katambora has a fine leaf, is early flowering and is less vigorous and palatable than other varieties. It is more persistent, forms a denser sward and shoots earlier in spring.

Callide is late flowering and takes a long time to respond in the spring. It



does best in fertile soils, is tall and late maturing. It has long awns and a long tuft of hairs at the awn base. Callide is more frost tender than Katambora and a smaller plant.

Pioneer is the original variety. It has a high proportion of stem and less leaf than the other varieties. It does not continue to grow into autumn and is less likely to respond to warmer winter weather. Topcut is derived from Pioneer Rhodes with a finer leaf and stem. It has an erect, uniform growth habit and is ideal for hay production. Finecut is a leafy Katambora type for better grazing and hay production. It forms a dense mat over the ground. Nemkat is resistant to nematodes.

Many seed companies now have their own selected lines of Rhodes grass under various trade names such as Mariner, Reclaimer and Toro.



## Sabi grass

*Urochloa mosambicensis*

Sabi grass is a perennial creeping grass that grows to 60 cm however it has variable size and habit depending on soil type and growing conditions.

There are four main types, three of which provide useful and palatable feed. Sabi grass tolerates hot and dry conditions but after frosting in winter it does not provide stand-over feed.

The favoured soil types for this grass depends on the variety but it will grow on many soil types from sands to light clays. Crude protein ranges from 10–20% in young growth.

This grass should not be confused with the annual grass *Urochloa panicoides* commonly found on cultivations during spring in the Burnett.

The Saraji variety is suited to sandy soils through to clay loams and areas of heavier rainfall. Nixon has good seedling survival, shorter growth habit and can persist in lower rainfall areas. Seed is readily available.



## Silk sorghum

*Sorghum arundinaceum*

Silk sorghum is an erect, robust, tussocky perennial with numerous tillers and thick short rhizomes that curve upwards to produce new shoots near the parental stool. Stems are solid and pithy, about 1 cm thick, sometimes reaching a height of 3–3.6 m. Leaves are 2.5–4 cm wide.

The seedhead has multiple branches that droop down as the seed ripens. Silk sorghum is suited to heavy, fertile soils and combines well with lucerne. It is easy to establish and will persist for one to five years, depending on soil management and rainfall.

Silk sorghum requires nitrogen fertiliser to grow and persist well. It is drought tolerant but does not handle flooding. It responds well to the onset of spring and soil temperatures above 15°C provide conditions for ideal growth. After long dry spells silk sorghum can be toxic due to high levels of prussic acid.

Silk is a hybrid between 'Krish' (a hybrid of Johnson grass (*S. halepense*) and *S. roxburghii*) and *S. arundinaceum*.

## Temperate grasses

There are a number of temperate climate grasses that do well in the Burnett during winter if there is sufficient rainfall or irrigation. Ryegrass, fescue, cocksfoot and phalaris are worth investigating, particularly if irrigation is available.



## Further information

### Department of Agriculture and Fisheries (DAF)

[www.daf.qld.gov.au](http://www.daf.qld.gov.au)

Customer Call Centre open normal business hours Monday to Friday (telephone 13 25 23 for the cost of a local call within Queensland; interstate callers 07 3404 6999) or email [CSCTeamLeader@daf.qld.gov.au](mailto:CSCTeamLeader@daf.qld.gov.au)

### FutureBeef



[www.futurebeef.com.au](http://www.futurebeef.com.au)

FutureBeef is a collaborative program for the northern Australia beef industry. The aim of the FutureBeef program is to support sustainable and profitable productivity gains for northern beef producers.

### Pastures Australia



[https://keys.lucidcentral.org/keys/v3/pastures/Getting\\_Started.htm](https://keys.lucidcentral.org/keys/v3/pastures/Getting_Started.htm)

Pastures Australia is an incorporated joint venture for investment in the genetic improvement, management and adoption of pasture plants across Australia. The partners are Australian Wool Innovation, the Grains Research & Development Corporation, Meat & Livestock Australia, Dairy Australia, and the Rural Research & Development Corporation.

The Pastures Australia *Pasture Picker* decision tool for farmers and advisors provides specific pasture recommendations for regions across Australia. Species selection can be refined to specific local conditions including climatic and soil variables.

### Pasture dieback

Many pastures in the Burnett, and Queensland-wide, have been affected by severe pasture dieback. The condition is characterised by the death of pasture in patches, often extending to whole paddocks. Pastures generally lose vigour and characteristic red leaves. Over time the condition results in death of pasture plants.

Pastures vary in susceptibility, with monocultures of creeping blue grass

and buffel grass most seriously affected. Mealybugs may be involved and the affected pasture may become covered in grey fungal material.

Research has found that pastures can recover, and where bare patches occur it is an opportunity to add legumes to the pasture. Dieback does not affect legumes. Research work continues into the cause of the condition but there have been no clear findings to date.

## Pasture photo-standards

[www.futurebeef.com.au](http://www.futurebeef.com.au)

Pasture photo-standards are an essential tool in good grazing land management. The *Pasture photo-standards* will assist in developing pasture budgets and dry season business management plans. The photo-standards and corresponding pasture yields for many of Queensland's common pasture communities can be found on the FutureBeef website. You will be able to search for standards according to region or by pasture type.

## Tropical forages

[www.tropicalforages.info](http://www.tropicalforages.info)

*Tropical Forages* is a powerful tool for selecting forage species suitable for local conditions in the tropics and subtropics. It is invaluable for agricultural researchers and extension officers involved in improving animal production.

*Tropical forages* allows you to speedily:

- identify forage species suitable for your climate, soils, production system and management via a selection tool built on LUCID™
- access comprehensive information on these species with details of adaptation, uses and management of forage species, cultivars and elite accessions
- view images of the plants and their use
- search a comprehensive database of scientific references with abstracts.



## Stocktake: balancing supply and demand

*Stocktake* is a paddock-scale land condition monitoring and management package. It has been developed to provide grazing land managers with a practical, systematic way to:

- assess land condition and long-term carrying capacity
- calculate short-term forage budgets.

The Stocktake GLM app now allows grazing land manager to undertake assessments and budgets in the paddock.

[www.stocktakeglm.com.au](http://www.stocktakeglm.com.au)

[www.futurebeef.com.au](http://www.futurebeef.com.au)

## Grazing Land Management

*Grazing Land Management* (GLM) workshops provide land managers with a practical and planned approach to improving productivity and sustainability. The workshops start by looking at grazing lands as ecosystems and how they function.

Held over three days, GLM workshops provide information that includes long-term stocking rate calculations, forage budgeting, land condition assessment and information on sown pastures, fire and weeds specific to the land types, climate and production systems of specific regions.

Check the FutureBeef website events page for details and registration for *Stocktake* and *Grazing Land Management* workshops.

[www.futurebeef.com.au](http://www.futurebeef.com.au)



# Species list

## Legumes

<i>Aeschynomene americana</i> and <i>A. falcata</i> (joint vetch).....	58
<i>Arachis</i> spp (forage peanuts).....	57
<i>Chamaecrista rotundifolia</i> (Wynn cassia).....	64
<i>Clitoria ternata</i> (butterfly pea).....	56
<i>Desmanthus virgatus</i> (Desmanthus).....	57
<i>Lablab purpureus</i> (lab lab).....	58
<i>Leucaena leucocephala</i> (leucaena).....	59
<i>Lotononis bainesii</i> (lotononis).....	59
<i>Lotus pedunculatus</i> (Maku lotus).....	61
<i>Macroptilium atropurpurem</i> (siratro).....	62
<i>Macroptilium bracteatum</i> (burgundy bean).....	55
<i>Medicago sativa</i> (lucerne).....	60
<i>Medicago</i> spp (annual medics).....	55
<i>Ornithopus compressus</i> (serradella).....	61
<i>Stylosanthes guianensis</i> var. <i>intermedia</i> (stylo, fine stem).....	63
<i>Stylosanthes scabra</i> (stylo, shrubby).....	63
<i>Stylosanthes seabrana</i> (stylo, Caatinga).....	62
<i>Trifolium</i> spp (clover).....	56
<i>Vicia villosa</i> ssp <i>dasycarpa</i> (vetch).....	64

## Grasses

<i>Bothriochloa bladhii</i> ssp. <i>glabra</i> (blue grass, Swann forest) .....	69
<i>Bothriochloa insculpta</i> (blue grass, creeping).....	68
<i>Bothriochloa pertusa</i> (blue grass, Indian) .....	70
<i>Brachiaria hybrids</i> .....	71
<i>Cenchrus ciliaris</i> (buffel grass).....	67
<i>Chloris gayana</i> (Rhodes grass) .....	75
<i>Dicanthium aristatum</i> (blue grass, Floren) .....	69
<i>Digitaria eriantha</i> (pangola grass).....	74
<i>Digitaria milanjiana</i> (digit grass).....	71
<i>Megathyrsus maximus</i> (was <i>Panicum maximum</i> ) (panics) .....	73
<i>Panicum coloratum</i> var. <i>makarikariense</i> (bambatsi) .....	67
<i>Paspalum dilatatum</i> (paspalum) .....	74
<i>Pennisetum clandestinum</i> (kikuyu) .....	72
<i>Setaria incrassata</i> (purple pigeon grass) .....	75
<i>Sorghum arundinaceum</i> (silk sorghum) .....	77
<i>Urochloa mosambicensis</i> (sabi grass).....	76
Temperate grasses .....	77



Supported by  
Natural Resources  
Investment Program

